

Insomnia in Pregnancy and Breastfeeding

MotherSafe - Royal Hospital for Women

Updated March 2022

Information in this leaflet is general in nature and should not take the place of advice from your healthcare provider. With every pregnancy there is a 3 to 5% risk of having a baby with a birth defect. Breastmilk provides optimum nutrition for babies and conveys many additional health benefits to mother and baby.

What is insomnia?

Insomnia is difficulty falling or staying asleep. It is common during pregnancy and in the months after birth when breastfeeding.^{1,2,3} For some women insomnia may be a symptom of depression or anxiety. If so, this should be assessed specifically with a healthcare professional.

Issues for pregnancy

In pregnancy, insomnia may happen for many reasons. These include changing hormones, nausea and vomiting, heartburn, needing to urinate frequently, leg cramps and restless legs, backaches, difficulty getting comfortable in bed from baby growing and moving, as well as stress, anxiety and depression.^{1,2,3}

Non-medical treatment

Good sleep hygiene, which involves improving habits around sleep is the main treatment for insomnia.

This includes measures such as ensuring regular times to go to sleep and wake up, keeping the bedroom cool and dark, avoiding electronic devices for an hour before going to bed and avoiding caffeine in the afternoon and evening (not advisable in pregnancy and breastfeeding anyway). Drinking water during the day, but stopping a few hours before going to bed will also reduce the need to pass urine throughout the night. Napping during the day should be avoided or kept to less than 20 minutes to allow for a better night-time sleep.^{1,2,3,4}

If there is no improvement with attention to sleep hygiene, consider seeing a doctor for further advice re possible non-medicinal therapy options (such as cognitive behavioural therapy and relaxation techniques) or temporary medicine options.^{1,2,3,4}

Medicines recommended

Medications are sometimes used in pregnancy to treat insomnia although this approach is generally **not** the mainstay of treatment.

A common group of medications used in pregnancy for insomnia are the sedating antihistamines such as doxylamine. Doxylamine has been widely used by pregnant women at all stages of pregnancy (including for treatment of morning sickness) and has not been associated with any increased risk to the baby. It is not habit forming.^{3,5}

Another class of medication used for insomnia is the benzodiazepine group. Temazepam is the most common benzodiazepine used for the treatment of insomnia. Continuous use of benzodiazepines is not advised as they are habit forming for the mother and may be associated with withdrawal symptoms in the baby after delivery.^{2,3,6} If this class of medicine is chosen, use the lowest effective dose for the shortest period of time. For advice regarding specific medications for treatment of insomnia, consult with your healthcare provider or call MotherSafe.

Melatonin, as a prescription medicine is sometimes used to treat insomnia. While there is little information about its use in pregnancy, there is no evidence of any harmful effects. However it may be preferable to use a medication with more evidence of safety in pregnancy.^{3,7} Homeopathic formulations of melatonin are available in Australia as an over-the-counter product. As these contain only minute amounts of melatonin, they would not be anticipated to cause problems for the baby but equally, they would not be expected to be effective in the treatment of insomnia.

Herbal medications are sometimes suggested to treat insomnia. They are generally unstudied in pregnancy so it is difficult to establish their safety for the baby, Furthermore, herbal products are not subject to the same degree of regulation as regular medicines, so the labelling may not accurately describe the actual ingredients (particularly if the product is not from a reputable supplier). Therefore, herbal medications are not recommended in pregnancy to treat insomnia.⁸

If there is underlying depression or anxiety, antidepressants may be considered and this should be discussed with a healthcare provider such as your GP or mental health professional. You can also call MotherSafe for advice about specific medications.

Other underlying conditions such as restless legs syndrome and heartburn may need to be treated specifically by your doctor.

Breastfeeding

Non-medical treatment

Sleep hygiene measures are the first line treatment of insomnia while breastfeeding (see advice above in: **Issues for pregnancy - Non-medical treatment**).

Medicines

It is not advisable to take any medication for insomnia while breastfeeding without medical advice or discussing with MotherSafe. This includes herbal remedies and over-the-counter medications. A particular concern about medication use for insomnia while breastfeeding is that some of the medication can pass into breastmilk and can lead to drowsiness in the baby. This is more likely when the baby is young (especially less than 2 months) and fully breastfed. If the decision is made to take medication, it is important to take the lowest effective dose for the shortest time period and ensure your baby is alert and waking for feeds. A short acting medication like temazepam⁹ is unlikely to be a problem particularly in an older baby on a one off or occasional basis and would be considered preferable to the over-the-counter antihistamine, doxylamine, for which there is no breastmilk excretion information available.¹⁰

Melatonin is relatively unstudied in breastfeeding but it is unlikely to be of concern in an older baby.¹¹

Herbal medications are unstudied in breastfeeding and are as likely to cause drowsiness and other side effects in the baby as regular medicines. Therefore they are not recommended for treatment of insomnia.

It is important to be very careful when handling or responding to your baby if you are taking a sleep inducing medicine due to the increased likelihood that you will be drowsy from the effects of the medicine.

Another concern with medications for insomnia is that some breastfeeding mothers may already be taking medications which may contribute to drowsiness and adding a sleep medicine may increase the likelihood of side effects in the baby. If you are taking other medications eg codeine or antidepressants, please discuss the addition of any new medication with an appropriate healthcare professional.¹²

Ask your midwife, doctor or pharmacist for the brand names of these medicines.

References

1. Miller MA, Mehta N, Clark-Bilodeau C, Bourjelly G. Sleep pharmacotherapy for common sleep disorders in pregnancy and lactation. Chest. 2020; 157(1):184-197
2. Therapeutic Guidelines (eTG March 2021 edition). Insomnia in the perinatal period. March 2021 West Melbourne. Accessed January 2022
3. Chaudry SK, Suser LC. Considerations in treating insomnia during pregnancy: a literature review. Psychosomatics 2018; 59:341-348
4. NPS MedicineWise|Consumer Info|Medical Info -How to sleep right. June 2017. Available at <https://www.nps.org.au/medical-info/consumer-info/how-to-sleep-right?c=tips-for-natural-sleep-8a0ccee3> Accessed December 2021
5. Rossi S (ed). Australian Medicines Handbook (internet). Doxylamine. January 2022. Adelaide. Australian Medicines Handbook Pty Ltd. Accessed March 2022
6. Therapeutic Guidelines (eTG March 2021 edition). Benzodiazepine use during pregnancy. March 2021 West Melbourne. Accessed January 2022
7. Therapeutic Guidelines (eTG March 2021 edition). Melatonin use during pregnancy. March 2021 West Melbourne. Accessed January 2022
8. Holst L, Wright D, Haavik S, Nordeng H. Safety and efficacy of herbal remedies in obstetrics-review and clinical implications. Midwifery 2011; 27:80-86
9. Lactmed; Dug and Lactation Database. Temazepam. National Library of Medicine. December 2021. Available at <https://www.ncbi.nlm.nih.gov/books/NBK501256/> Accessed March 2022

10. Lactmed; Drug and Lactation Database. Doxylamine. National Library of Medicine. December 2021. Available at <https://www.ncbi.nlm.nih.gov/books/NBK500620/> Accessed March 2022
11. Lactmed; Drug and Lactation Database. Melatonin. National Library of Medicine. December 2021. Available at <https://www.ncbi.nlm.nih.gov/books/NBK501863/> Accessed March 2022
12. Chow KC and Koren G. Sedating drugs and breastfeeding. Canadian Family Physician March 2015; 61:241-243

Other resources

MotherToBaby. When counting sheep doesn't work: Insomnia and pregnancy. OTIS. Organization of Teratology Information Specialists. December 2016. Available at <https://mothertobaby.org/baby-blog/when-counting-sheep-doesnt-work-insomnia-and-pregnancy/>

Australian Breastfeeding Association. How to cope with broken sleep. April 2021. Available at <https://www.breastfeeding.asn.au/bfinfo/how-cope-broken-sleep>



NSW Medications in Pregnancy & Breastfeeding Service

For more information call MotherSafe: NSW Medications in Pregnancy and Breastfeeding Service on 9382 6539 (Sydney Metropolitan Area) or 1800 647 848 (Non-Metropolitan Area) Monday -Friday 9am-5pm (excluding public holidays)