

SESLHD GUIDELINE COVER SHEET



NAME OF DOCUMENT	Constipation - Prevention and Management
TYPE OF DOCUMENT	GUIDELINE
DOCUMENT NUMBER	SESLHDGL/098
DATE OF PUBLICATION	December 2025
RISK RATING	Medium
LEVEL OF EVIDENCE	National Safety and Quality Health Service Standards: Standard 1 – Clinical Governance Standard 2 – Partnering with Consumers Standard 4 – Medication Safety Standard 5 – Comprehensive Care Standard 6 – Communicating for Safety NSW Health PD2022_032 - Medication Handling in NSW Public Health Facilities SESLHD Nurse Initiated Protocols
REVIEW DATE	December 2028
FORMER REFERENCE(S)	SESLHDGL/098 – Bowel Management
EXECUTIVE SPONSOR	Clinical Stream Director, Medicine
AUTHOR	SESLHD Bowel Assessment Working Party
POSITION RESPONSIBLE FOR DOCUMENT	Clinical Stream Manager, Medicine carolyn.smith1@health.nsw.gov.au
FUNCTIONAL GROUP(S)	Aged Care and Rehabilitation Cancer and Palliative Care Services Medicine Surgery, Perioperative and Anaesthetic
KEY TERMS	Constipation, Digital Rectal Examination (DRE), Enema Administration
SUMMARY	To provide guidance and advise on the procedures for the successful assessment, management and prevention of constipation in adult inpatients to ensure safe and effective patient management.

THIS DOCUMENT IS A GUIDE FOR BEST PRACTICE
This Guideline is intellectual property of South Eastern Sydney Local Health District.
Guideline content cannot be duplicated.

Feedback about this document can be sent to SESLHD-Policy@health.nsw.gov.au

Constipation – Prevention and Management

Section 1 - Background	3
Section 2 - Principles	4
Expected Outcomes of a Program to Prevent/Manage Constipation	4
Aims of a Program to Prevent/Manage Constipation	4
Principles of Effective Prevention/Management of Constipation	4
Risk Factors for Constipation	5
Section 3 - Definitions	6
Section 4 - Responsibilities	8
Nurse /Midwifery Unit Managers (NUM/MUM) / Department Managers are responsible for:.....	8
Medical Officers are responsible for:.....	8
Nurse Practitioners (NP) are responsible for:	8
Registered Nurses (RN) / Midwives (RM) are responsible for:	8
Enrolled Nurses (EN) are responsible for:	8
Assistants in Nursing (AIN) are responsible for:	9
Section 5 - Assessment and Documentation on Admission to Hospital or Community Service – All Patients	10
Section 6 - Constipation Management	11
Prevention and Treatment for All Patients	11
Low Risk of Constipation.....	11
Patients with Risk Factors	11
High Risk of Constipation.....	11
Opioids	11
Section 7 - Discharge Planning for Prevention/Management of Constipation	13
Aims and expected outcomes	13
Discharge management and follow-up	13
Section 8 - Laxatives (Recommended)	14
Section 9 - Laxatives (Other).....	17
Section 10 - Dosage Options for Osmotic Laxatives Including Enema Recommendations for Faecal Impaction. 19	
Dosage for prevention of constipation.....	19
Dosage for Constipated Patients	19
Dosage for Faecal impaction (Option 1 – 4, as per treating team).....	19
Administration via an Enteral Feeding Device	19
Section 11 - Documentation of Admission Assessment and Physical Examination	20
Current history.....	20
Findings of abdominal assessment.....	20
Findings of Digital Rectal Examination (DRE) if applicable – refer to Appendix E.	20
Section 12 - References	21
Section 13 - Version and Approval History.....	22
Appendix A: Abdominal Assessment	23
Appendix B: Digital Rectal Examination (DRE).....	25
Limitations for Practice	25
Indications for Digital Rectal Examination.....	25
Clinical history to be obtained prior to digital rectal examination	25
Precautions	25
Contraindications.....	26
Preparation for procedure	26
Procedure.....	26
Appendix C: Procedure for Administration of an Enema.....	28
1. Fleet Enema	28
2. Olive Oil Enema	29
Appendix D: Procedure for Manual Disimpaction	30
Appendix E: Interventions Related to DRE Findings.....	31

Section 1 - Background

Successful management of constipation and altered bowel habits depends on the assessment and identification of the underlying cause and/or other contributory factors.

Prevention of constipation is imperative and a direct nursing responsibility.

Early and accurate assessment, followed by preventative measures can eliminate the need for more invasive treatments.

All information on bowel assessment and management should be clearly and consistently documented on a daily basis. This information should be handed over across all shifts and on inter-ward and inter-hospital transfers.

RESOURCE PERSONNEL

The ward based Clinical Nurse Specialists (CNS) and Clinical Nurse Educators (CNE) should be contacted to assist with bowel management in the first instance

Other Specialist or Clinical Nurse Consultant (CNC) teams and allied health professionals may be contacted for further consultation during business hours.

The Sutherland Hospital	
Ward / Unit Clinical Nurse Educators / Specialist	Palliative Care
Admitting Medical Team	Continence
	Aged Care
St George Hospital	
Ward / Unit Clinical Nurse Educators/ Specialist	Palliative Care
Admitting Medical Team	Continence
	Aged Care
	Anorectal
Prince of Wales Hospital	
Ward / Unit Clinical Nurse Educators/ Specialist	Palliative Care
Admitting Medical Team	Continence
	Aged Care
	Anorectal
Sydney Hospital and Sydney Eye Hospital	
Admitting Medical Team	
Nurse Manager, Clinical Practice & Development, and Leadership Capabilities	

Section 2 - Principles

EXCLUSIONS

Patients with the following conditions will NOT have Digital Rectal Examination (DRE) or administration of per rectum aperients before consultation with the treating medical team.

- Patients requiring serial abdominal assessment post trauma
- Existing bowel pathology e.g.: inflammatory bowel disease, Crohn's disease, ulcerative colitis, toxic megacolon
- Following bowel or rectal surgery or patients with rectal bleeding:
- Patients with ileostomy or jejunostomy
- Patients unable to tolerate enteral feeding/oral diet:
- Patients with raised intracranial pressure - consult the treating intensivist or neurosurgeon before performing PR examination or aperient administration
- Neutropenic patients
- Patients with a Platelet count of less than 30 (Confirmation of platelet count would need to be obtained from Haematology)
- Spinal injury patients will be managed according to local policy and ACI regime

Expected Outcomes of a Program to Prevent/Manage Constipation

- Patient establishes a pattern of elimination consistent with his or her norm and is comfortable in relation to elimination
- Nursing strategies are implemented to modify or restore to normal the patient's pattern of elimination
- Nursing strategies are evaluated daily, altered according to the patient's elimination pattern and level of comfort and documented in the clinical notes/bedside handover tool.

Aims of a Program to Prevent/Manage Constipation

- Achieve regular effective and efficient emptying (evacuation) of the colon avoiding intermittent treatment for faecal overloading or impaction
- Prevent constipation, faecal impaction and resultant faecal incontinence
- Effectively treat impaction and overload and return bowel function to normal.

Principles of Effective Prevention/Management of Constipation

- Patients using bulking agents must be able to tolerate the increased fluid intake required for use of these agents, otherwise they are at risk of obstruction or impaction.
- Patients using bulking agents must have normal colonic transit i.e. **NO** underlying neurogenic conditions such as Parkinson's disease, slow transit constipation, MS, motor neurone disease.
- When laxatives are required select a recommended laxative ([Section 6](#)) and avoid using those which must be used with caution ([Section 7](#)).

- All patients with chronic bowel management problems should be referred to a specialist nurse for assessment.
- If bowels do not open for 3 days and recommended interventions have been implemented, the patient requires full assessment for possible faecal impaction/overload/obstruction, including medical review and possible abdominal x-ray +/- DRE.

Risk Factors for Constipation

- Immobility
- Spinal injury
- Lower back injury / surgery particularly S1 – S4
- Neurological diseases e.g. Parkinson's disease, MS, motor neurone disease and dementia.
- Delirium
- Cognitive/psychiatric deficits
- Inadequate dietary intake and low fibre diets
- Reduced fluid intake/dehydration
- Modified diets such as puree and thickened fluids
- Communication deficits including aphasia and CALD
- Pain e.g. Anal fissures or haemorrhoids, recent surgery
- Patients who have regular laxatives, now ceased due to hospitalisation
- Medications e.g.: [Opioids](#), [drugs with anticholinergic effects](#), antidepressants, aluminium-containing antacids, verapamil, 3HT₃-receptor antagonists (e.g. ondansetron), and calcium or iron supplements
- Physiological and psychological barriers such as lack of privacy, bedpans, shared bathroom.

Section 3 - Definitions

Definition: Ballooned Rectum

- Rectum feels dilated on examination.
- A ballooned (dilated) rectum is indicative of gross constipation
- An empty ballooned rectum may indicate faeces high up in the bowel, or it may be a sign of bowel obstruction
- Consider abdominal X-Ray.

Definition: Bulking Agents (Fibre)

- A form of fibre that is used in the management of constipation
- Aims to absorb water into the digestive tract to stimulate the muscles in the bowel to encourage regular bowel movements
- Bulk the stool and make it easier and softer to pass.

Definition: Collapsed Rectum

- The rectal wall should be easily felt collapsed around the examining finger
- This usually indicates a functioning bowel.

Definition: Constipation

- Constipation refers to reduced stool frequency and/or difficulty passing stool (e.g. straining, incomplete evacuation, hard/lumpy stools, prolonged time to pass stool, need for manual manoeuvres). It is a highly subjective symptom with normal bowel habits varying between individuals
- Defining constipation must involve measurable, objective symptoms including frequency of defecation and stool characteristics, as well as the patient's perception of constipation relating to ease of defecation and associated level of discomfort
- Definition by National Institute of Diabetes and Digestive and Kidney Diseases is bowel movement less than 3 times a week, that is hard, dry and/or lumpy and the evacuation is difficult
- Although constipation is exacerbated by **opioids** and anticholinergic drugs, the cause of constipation is usually multifactorial
- The consequences of unmanaged constipation include abdominal pain, bloating, nausea, overflow incontinence, tenesmus, faecal impaction, urinary retention and occasionally bowel obstruction and colonic perforation, delirium; all of which can profoundly affect the quality of life.

Definition: Digital Rectal Examination (DRE)

- A DRE is the examination of the rectum with a gloved and lubricated finger for the purpose of:
 - Establishing the presence, amount and consistency of faecal matter in the bowel
 - Assess for haemorrhoids, fissures, abscesses or tumour masses
 - Assessing the need for rectal medication or digital removal of faeces in extreme cases of faecal impaction
 - Gauging anal sphincter function and tone and rectal sensation.

Definition: Enema

- An enema is the introduction of fluid through the anus into the rectum and colon
- A retention enema is fluid that is introduced into the rectum to be retained.

Definition: Faecal Impaction

- A solid immobile bulk of faeces that develops in the rectum as a result of chronic constipation
- May also suffer from faecal incontinence to Bristol stool type 6-7 due to overflow
- Faecal impaction must be clinically diagnosed
- Abdominal X-ray **may** be required to determine extent of impaction. (*A plain abdominal X-ray is of limited utility in diagnosis of constipation and is mainly used to exclude bowel obstruction*).

Definition: Faecal Loading

- A large volume of stool hard or soft that accumulates in the rectum and sigmoid colon; often presents as faecal incontinence
- May also suffer from faecal incontinence to Bristol stool type 6-7 due to overflow.

Definition: Laxatives (Aperients)

- Medication commonly used to treat constipation and classified by the mode of action into categories of; stimulant laxatives, faecal softeners, osmotic laxatives and rectal suppositories.

Definition: Manual Disimpaction

- Manual disimpaction is the digital removal of stool from the rectum using a gloved and lubricated finger. Refer to [Appendix D](#).

Section 4 - Responsibilities

Nurse /Midwifery Unit Managers (NUM/MUM) / Department Managers are responsible for:

- Ensuring compliance with this guideline in their unit department.

Medical Officers are responsible for:

- Undertaking regular review of patients based on clinical assessment and when alerted to an issue by nursing staff.
- Documentation of clinical assessment findings and a clear management plan.
- The prescribing of appropriate bowel medications and regular review of the response to those medications.

Nurse Practitioners (NP) are responsible for:

- Working within their individual scope of practice.
 - Undertaking regular review of patients based on clinical assessment
 - Documentation of clinical assessment findings and a clear management plan.
 - The prescribing of appropriate bowel medications and regular review of the response to those medications.

Registered Nurses (RN) / Midwives (RM) are responsible for:

- Daily assessment and evaluation of the patients' bowel habits to identify altered bowel habits according to the patient elimination pattern and level of comfort.
- Documenting findings as per [Section 8](#) in the patients eMR record & eMR bowel chart.
- Implementing strategies to modify or restore the patients' normal pattern on elimination, including the administration of nurse initiated and prescribed medication as appropriate.
- Escalate failure to respond to interventions to the treating team and N/MUM.

Enrolled Nurses (EN) are responsible for:

- Daily assessment and evaluation of the patients' bowel habits to identify altered bowel habits according to the patient elimination pattern and level of comfort.
- Documenting findings as per [Section 8](#) and report findings to the Registered Nurse.
- Implement strategies to modify or restore the patients' normal pattern on elimination, including the administration of nurse initiated and prescribed medication within scope of practice.

NOTE: Only ENs without notation may administer medications as per [SESLHDPD/160 - Medication: Administration by Enrolled Nurses](#) and [SESLHD Nurse Initiated Protocols](#).

Assistants in Nursing (AIN) are responsible for:

- Daily assessment and evaluation of the patients' bowel habits to identify altered bowel habits according to the patient elimination pattern and level of comfort
- Documenting findings as per [Section 8](#) and report findings to the Registered Nurse
- Liaise with Medical and Nursing/Midwifery staff regarding nutritional management
- Document assessment and intervention in the eMR medical record.

Section 5 – Assessment and Documentation on Admission to Hospital or Community Service – All Patients

- On admission initial assessment is completed in the Elimination Section on Adult Admission Assessment (AAA). It is important to gather additional information if there are concerns that the patient is at risk of altered elimination. Appropriate assessment should include interviewing the patient/significant other to obtain:
 - Previous bowel history
 - Usual patterns/continence status
 - Existing bowel status
 - Laxative use
- Identify known risk factors (Refer to [Section 2](#) page 5) for constipation and resultant overflow/faecal incontinence
- DAILY documentation in eMR is the minimum standard of bowel status
- Documentation on the eMR Bowel chart is required for all patients – especially those at risk of constipation and consider need for a fluid balance chart & fluid balance summary chart
- Appropriate assessment strategies are supported by documentation prior to implementing prevention and/or treatment of constipation or faecal impaction
- Evaluation and outcomes related to assessment and management regime must be included in the clinical notes where a problem or potential problem exists
- Documentation - refer to [Section 8](#).

Physical Examination

Undertake an abdominal examination if the patient:

- Has failed to pass stool (as per Bristol Stool Form Scale on the Bowel Management Chart in eMR) after 3 consecutive days
- Is confused and unable to give a history (family, carer or facility cannot provide the information)
- Abdominal Assessment - inspect, auscultate, percuss and palpate all abdominal quadrants if trained to do. See [Appendix A](#).

NOTE:

- Physical examination may include a DRE (refer to [Appendix B](#)) in consultation with the medical team. An abdominal X-ray may also be required to assess for impaction.

Section 6 - Constipation Management

Prevention and Treatment for All Patients

- Conduct assessment and physical examination if indicated
- Document findings in the patients eMR record & eMR bowel chart
- Monitor and document bowel status daily as a minimum
- Ensure the patient's regular laxatives (if applicable) are prescribed and administered
- Implement recommendations according to risk status

Low Risk of Constipation

Patients with minimal or no risk factors for constipation as per [Section 2](#) Risk Factors.

If bowels not opened for 2 days, commence a low dose of oral aperients

Recommended agents are (also refer to [Section 6](#))

- Macrogol 3350 (Osmolax 1-2 scoops daily), OR
- Macrogol 3350 with electrolytes (Movicol 1-3 sachets daily)
- Consider Psyllium Husk powder

Patients with Risk Factors

- If bowels not opened for 2 days, discuss management plan with the patient.
- Consult a ward CNS/CNE then treating medical team for advice (for additional resource personnel refer to [Section 1](#)).
- Suggest DRE in appropriate patients and manage as per flow chart "Interventions related to DRE findings" – refer to [Appendix E](#).
- If patient unwilling to consent to DRE, continue oral aperients as above.
- If bowels not opened for 3 days, patient requires a DRE in appropriate patients and further management as per flow chart in [Appendix E](#).
- Reassess risk and titrate aperients daily according to bowel action.

High Risk of Constipation

Patients at moderate to high risk of constipation (refer to [Section 2](#) page 5)

- If no regular aperients taken at home, or aperients are ineffective, commence aperient according to individual patient assessment and risk factors (refer to [Section 6](#)).
- If bowels not opened for 2 days, discuss management plan with patient.
- Perform DRE in appropriate patients and manage as per flow chart [Appendix E](#).
- Reassess risk and titrate aperients daily according to bowel action.

Opioids

Patients receiving opioid medications are at particularly high risk of constipation.

- Docusate & Senna are available in combination (Coloxyl & Senna[®]) and are the aperients of choice as they counteract the effect of morphine on the gut with dual stimulant and softener action
- Recommended dose is 2 tablets at night.
- If not effective, refer to the treating medical team. The medical team can increase the dose to up to two tablets twice a day.

Section 6 Constipation Management

- Refer to [SESLHDGL/111 - Prevention and Management of Opioid Induced Constipation in advanced palliative care patients](#) for management plan if on Opioids.

Section 7 - Discharge Planning for Prevention/Management of Constipation

For specific advice on discharge management refer to the relevant medical recommendations as per patients discharge summary.

Aims and expected outcomes

- All patients to be discharged with an established elimination pattern OR appropriate bowel management strategies are identified and patient education attended pre discharge
- Return to normal premorbid bowel habit on discharge
- Management strategies for alterations to elimination pattern are instigated.
- Review patient history of bowel activity during admission
- Review patient's long-term history
- Identify if intervention required post discharge
- Identify action required and timeframe
- Document in Discharge Summary and advise for review/follow up with GP, if further management is required.

Discharge management and follow-up

- Establish bowel pattern both premorbid and current
- Identify possible causes of altered elimination pattern e.g. poor oral intake, fluid restriction, lack of exercise, medications
- If dietary causes identified, refer to dietician as inpatient or outpatient.
- Document in Discharge Summary and advise for review/follow up with GP, if further management is required.

Section 8 – Laxatives (Recommended)

Refer to [SESLHD Nurse Initiated Protocols](#)

BULKING AGENTS	TYPE	ACTION	COMMENTS
Psyllium (Metamucil) Wheat Dextrin (Benefiber)	Powder	Bulking agent / fibre supplement	Increases the faecal mass which stimulates peristalsis and produces bulkier, softer stools. Needs adequate fluid intake
SOFTENER	TYPE	ACTION	COMMENTS
Sodium Docusate (Coloxyl)	Tablets	Faecal softener	Suitable for all patients as a first line treatment for constipation
Sodium Poloxamer (Coloxyl)	Drops (infant)	Faecal softener	On formulary however, POWH does not have in stock
OSMOTICS	TYPE	ACTION	COMMENTS
Macrogol 3350 (Osmolax) Macrogol 3350 with electrolytes (Movicol) – see also 4.17	Powder	Delivers water to the colon, softens stool and increases peristalsis. Products <i>with electrolytes</i> are iso-osmotic, which minimise loss of sodium, potassium and water.	Caution in fluid restriction and patients who may not tolerate fluid intake: Each level scoop of Osmolax will need to be mixed with 240mL of water. Patients need to be able to tolerate drinking 125 mL water or juice per sachet of Movicol. Patients can sip on this over several hours if volume is too large to drink at once. Can have up to 8 sachets of products <i>with electrolytes</i> (i.e. Movicol) per day for faecal impaction. (If concern for faecal impaction, consult medical officer.) Not suitable for patients certain modified diets e.g. Level 3 thickened fluids.
Lactulose (Actilax, Dulose)	Syrup	Water is drawn into the bowel, increasing the fluid volume	Requires a high fluid intake and consistent usage over 1-3 days for onset. Can cause fluid and electrolyte imbalances,

Section 8
Laxatives (Recommended)

		softening the stool and increasing peristalsis	diarrhoea and abdominal cramps. Is not suitable for the elderly in large doses. Is suitable for diabetics for short term use, however other laxatives preferred. Higher doses are used for hepatic encephalopathy.
STIMULANTS	TYPE	ACTION	COMMENTS
Bisacodyl (Dulcolax®)	Tablets	Increases pressure waves in colon and stimulates emptying.	Maintain adequate water intake. Can cause cramps and bloating. Stimulant laxative use is contraindicated in intestinal obstruction, acute abdominal conditions and inflammatory bowel conditions.
Docusate & Senna (Coloxyl & Senna ®)	Tablets	Faecal softener and peristaltic stimulant	Maintain adequate water intake. Can cause cramps and bloating. Stimulant laxative use is contraindicated in intestinal obstruction, acute abdominal conditions and inflammatory bowel conditions.
Sodium picosulfate (Dulcolax®)	Drops	Stimulates peristalsis. May promote accumulation of water and electrolytes into the colonic lumen.	Maintain adequate water intake. Can cause fluid and electrolyte imbalances. Can cause cramps and bloating. Stimulant laxative use is contraindicated in intestinal obstruction, acute abdominal conditions and inflammatory bowel conditions. Approved for use where solid oral dose forms of stimulant laxatives are not suitable.
RECTAL LAXATIVES	TYPE	ACTION	COMMENTS
Glycerol suppository	Suppository	Glycerol is a mild irritant. It works by encouraging muscles in the anus to contract. Glycerol lubricates and softens stools.	Lie patient in left, lateral recumbent position. Suitable for all age groups as first line management Should be inserted rectally into the stool.

Section 8
Laxatives (Recommended)

Bisacodyl suppository (Dulcolax suppository)	Suppository	Stimulate the bowel lining to increase peristalsis	Lie patient in left, lateral recumbent position. Suitable to be given in combination with glycerol suppositories. Should be placed against the rectal wall as it is absorbed through the rectal mucosa. May cause cramping.
---	-------------	--	---

Section 9 – Laxatives (Other)

THESE LAXATIVES MUST BE PRESCRIBED BY A MEDICAL OFFICER and/or NURSE PRACTITIONER - working within their individual scope of practice.			
Fleet Enema (Sodium Phosphate)	Enema	Draws and retains water into the bowel to soften stool and make it easier to pass.	<p>Lie patient in left, lateral recumbent position. A DRE must be performed prior to insertion to ensure no contraindications for administering the fleet enema.</p> <p>**At POWH the initial DRE must be performed by a medical officer**</p> <p>Use with caution in patients with impaired renal function, heart disease or pre-existing electrolyte disturbances. Also, patients on calcium channel blockers, diuretics or other medications that may affect electrolyte levels.</p> <p>Repeated dosage may result in hyperphosphataemia, hypokalaemia, hypocalcaemia, hypernatraemia and acidosis may occur. Adequate hydration via oral fluids should be given, to reduce the risk of dehydration and electrolyte disturbance where required.</p> <p>An irrigation / enema cone must be used for the administration of a Fleet enema at POW (refer to Appendix C). Administering the fleet enema without caution may cause adverse outcomes including bowel perforation.</p> <p>Advise patient to remain in position and hold contents of enema in rectum for at least 10 minutes, or as long as possible.</p> <p>If after the enema solution is administered there is no return of liquid, contact a doctor immediately as dehydration could occur.</p> <p>**It is not necessary to empty the bottle completely and do not force all contents in.</p> <p>The outcome of the examination needs to be documented on eMR medical progress note clearly.</p>
Microlax Enema	Enema	Acts to increase intestinal water secretion and stimulate peristalsis. Each 5 mL enema contains sodium citrate 450 mg, sodium lauryl	<p>Lie patient in left, lateral recumbent position.</p> <p>A DRE must be performed prior to insertion.</p> <p>Gently insert the lubricated enema tip into the rectum. If tip does not insert freely</p>

Section 9
Laxatives (Other)

		<p>sulfoacetate 45 mg, sorbitol 3.125 g.</p>	<p>or patient complains of pain, stop immediately. Seek medical advice before continuing.</p> <p>Advise patient to remain in position and hold contents of enema in rectum for at least 10 minutes, or as long as possible.</p>
Bisacodyl enema	Enema	<p>Stimulant laxative.</p> <p>Act by direct stimulation of nerve endings in colonic mucosa to increase intestinal motility. May also cause accumulation of water and electrolytes in the colonic lumen.</p> <p>Indicated for constipation in palliative care, where there is an empty rectum.</p>	<p>Lie patient in left, lateral recumbent position.</p> <p>A DRE must be performed prior to insertion.</p> <p>Shake well before use.</p> <p>Carefully insert the applicator nozzle tip into the rectum, then press on the end of the plastic tube to squeeze the liquid out gently.</p> <p>Remain lying down for a couple of minutes before sitting or standing up.</p> <p>Adverse effects - diarrhoea, fluid and electrolyte imbalance (with prolonged use or excessive doses), rectal irritation or burning, abdominal discomfort, cramps, nausea.</p>
Olive Oil Enema	Oil	<p>Rectal lubricant.</p> <p>Considered for palliative care patients to avoid manual disimpaction.</p>	<p>Lie patient in left, lateral recumbent position.</p> <p>A DRE must be performed prior to insertion.</p> <p>Olive oil (100-150mL) is retained in the rectum overnight to soften hard or impacted faeces to allow easier evacuation. Do not expect immediate results. The enema may need to be repeated daily before evacuation is achieved. Efficacy depends on the patient's ability to retain the oil. This is usually administered in conjunction with oral aperients or followed by stimulant suppository 12 hours after oil administration.</p>

Section 10
Dosage Options for Osmotic Laxatives Including Enema
Recommendations for Faecal Impaction)

Section 10 - Dosage Options for Osmotic Laxatives Including Enema Recommendations for Faecal Impaction.

Dosage for prevention of constipation

Movicol: Dissolve the contents of 1 sachet in 125mL of water and have the patient drink the mixture.

Osmolax: Dissolve 1 level scoop in 240mL of water.

Dosage for Constipated Patients

- Commence with 1 sachet of Movicol or 1 scoop of Osmolax per day.
 - Opiate induced constipation may need a higher dose.
- Adjust the dose upwards by 1 sachet per day up to a maximum of 3 sachets of Movicol (or 2 scoops of Osmolax) per day until a bowel movement is achieved.
- Once bowels are moving, use the Bristol Stool scale to titrate Movicol dose to achieve a Type 3 or Type 4 stool.

Dosage for Faecal impaction (Option 1 – 4, as per treating team)

- Faecal impaction must be clinically diagnosed by abdominal imaging.
- Abdomen X-ray may be necessary to determine the amount of impaction +/- a DRE.
- Option 1: Oral – Movicol, 4 to 8 sachets, each sachet dissolved in 125mL water, daily up to 3 days.
- Option 2: Suppository – Glycerol +/- Bisacodyl, rectally as a single dose.
- Option 3: Enema - Microlax® rectally, as a single dose to clear the distal bowel if suppository ineffective. If ineffective, Fleet® enema STAT. Refer to [Section 7 – Laxatives \(Other\)](#) and [Appendix C](#).
- Option 4: If option 1-3 are ineffective, escalate to the treating team. Consider manual disimpaction. Refer to [Appendix D](#).
- In Palliative Care, Dulcolax® drops are often used when solid oral stimulant laxatives are unsuitable, and an olive oil enema may be considered to help avoid the need for manual disimpaction.
- Encourage oral fluids and ambulation if patient is able. If oral fluids are limited or patient on a fluid restriction, suggest IV hydration.

Administration via an Enteral Feeding Device

- DO NOT use intravenous infusion devices for administering medications or fluid via the enteral route.
- Administer immediately after preparation by a manual bolus syringe method and flush with water following administration before the residue thickens in the tube.
- For administration via jejunostomy administer in 50 mL increments.
- Ensure the patient has been referred to a dietician so that the feeding regimen can be reviewed and changed if clinically indicated.
- Consult pharmacy for review of medication administration via an enteral device.

Section 11 - Documentation of Admission Assessment and Physical Examination

Following admission assessment and physical examination document the following patients' eMR record.

Current history

- Normal activities to maintain bowel function (e.g. what works for the patient)
- Any history of bowel surgery / conditions
- Changes in bowel habits
- Changes in amount of flatus passed
- Feeling of incomplete emptying
- Straining and/or inability to pass stool
- Type of stool, as per the Bristol Stool Form Chart
- If recent illness what was their normal bowel pattern prior to illness
- Identify aperient use – type, frequency and length of time used
- Individual preferences and special toileting needs e.g. raised toilet seat, foot stool etc
- Any pain and or blood with defaecation
- Presence of faecal incontinence
- Risk factors for constipation ([Section 2](#) page 5).

Findings of abdominal assessment

- Presence/absence of abdominal distention or bloating
- Presence/absence of bowel sounds.

Findings of Digital Rectal Examination (DRE) if applicable – refer to [Appendix E](#).

Daily assessment and evaluation of the patients' bowel habits to identify altered bowel habits according to the patient elimination pattern and level of comfort.

- DAILY documentation in eMR is the minimum standard of bowel status
- Documentation in the eMR Bowel Chart is required for all patients – especially those at risk of constipation and consider need for a fluid balance chart & fluid balance summary chart

Section 12 - References

1. Alarcon de la Lastra C, Barranco MD, Motilva V, Hurerias JM. (2001). Mediterranean Diet and Health: biological importance of olive oil. *Current Pharmaceutical Design*; (10) 933-950
2. Calvary Health Care Sydney Policy Number: 35 -: Bowel Management: Palliative Care
3. Candy B, Jones L, Larkin PJ, Vickerstaff V, Tookman A, Stone P (2015), Laxatives for the management of constipation in people receiving palliative care. *The Cochrane Database of Systematic Reviews*.
4. Coggrave M, Norton C, Cody JD (2014), Management of faecal incontinence and constipation in adults with central neurological diseases. *The Cochrane Database of Systematic Reviews*.
5. De Schryver, AM, Keulemans YC, Peters HP, Akkermans LM, Smout AJ, De Vries WR, van Berge-Henegouwen GP. (2005). Effects of regular physical exercise on defaecation pattern. *Scand J Gastroenterol*.
6. Jackson R, Cheng P, Moreman S, Davey, N & Owen, L (2016), "The constipation conundrum": Improving recognition of constipation on a gastroenterology ward. *BMJ Quality Improvement Reports*.
7. Kumar, L, Barker, C & Emmanuel, A (2014), Opioid-induced constipation: pathophysiology, clinical consequences and management. *Gastroenterology Research and Practice*, vol 2014
8. Lee-Robichaud H, Thomas K, Morgan J, Nelson RL (2010), Lactulose versus Polyethylene Glycol for Chronic Constipation. *The Cochrane Database of Systematic Reviews*
9. Lizarondo, L (2018), Constipation in Hospitalized Patients: Management. *The Joanna Briggs Institute EBP Database*.
10. Long Khanh-Dao Le, B.Pharm, (2018). Constipation Management (Older Person). *The Joanna Briggs Institute EBP Database*.
11. O'Laughlen, M 2009, Making sense of abdominal assessment, *Nursing Made Incredibly Easy*, vol. 7, no 5, pp. 15-19, Lippincott Williams & Wilkins, Inc.
12. Peate, I. (2003). *Nursing role in the management of constipation: use of laxatives*. *British Journal of Nursing* 12 (19) pp 1130 -36.
13. Powell, M., Rigby, D. (2000). *Management of bowel dysfunction and evacuation difficulties*. *Nursing Standard* 14 (4) pp 47-51.
14. Royal College of Nursing (2019) *Bowel Care Management of lower bowel dysfunction including Digital Rectal Examination and Digital Removal of Faeces*. London. RCN
15. Richmond, J. (2003). Prevention of Constipation through risk management. *Nursing Standard*, 17(16), 39-48.
16. Cherny N, Fallon M, Kaasa S, Portenoy R & Currow D (eds) 2015 *Oxford Textbook of Palliative Medicine* (5th Ed), Oxford University Press, Oxford. pp 675-685.
17. Wisten, A & Messner, T. (2005) Fruit and Fibre in the Prevention of Constipation. *Scandinavian Journal of Caring Sciences* Vol 19 p71.
18. <https://www.nursingtimes.net/clinical-archive/long-term-conditions/management-of-ascites-in-patients-with-liver-disease-18-09-2018/>

Section 13 - Version and Approval History

Date	Version	Version and approval notes
January 2022	DRAFT	Initial draft. Published on Draft for Comments page.
June 2022	DRAFT	Final version approved by Executive Sponsor. To be tabled at Quality Use of Medicines Committee and Clinical and Quality Council.
July 2022	DRAFT	Endorsed by the Quality Use of Medicines Committee with minor amendments.
September 2022	1	Approved at the Clinical and Quality Council meeting.
9 December 2025	1.1	<p>Minor review. Title changed.</p> <p>Section 1 - Resource Personnel; added information for SSEH</p> <p>Section 3 - Faecal Loading definition amended.</p> <p>Section 9 - added Bisacodyl enema</p> <p>Section 9 - NURSE PRACTITIONER - working within their individual scope of practice, was included.</p> <p>Section 10 - Updated options, dosage options for osmotic laxatives Including enema recommendations for faecal impaction.</p> <p>Dosage for faecal impaction (Option 1 – 4, as per treating team)</p> <ul style="list-style-type: none"> Option 1: Oral – Movicol, 4 to 8 sachets, each sachet dissolved in 125mL water, daily up to 3 days. Option 2: Suppository – Glycerol +/- Bisacodyl, rectally as a single dose. Option 3: Enema - Microlax® rectally, as a single dose to clear the distal bowel if suppository ineffective. If ineffective, Fleet® enema STAT. Refer to Section 9 – Laxatives (Other) and Appendix C. Option 4: If option 1-3 are ineffective escalate to the treating team. Consider manual disimpaction Appendix D. <p>Removed Pico prep as an option for faecal impaction.</p> <p>Added the following to Section 11 - Daily assessment and evaluation of the patients' bowel habits to identify altered bowel habits according to the patient elimination pattern and level of comfort.</p> <ul style="list-style-type: none"> DAILY documentation in eMR is the minimum standard of bowel status Documentation in the eMR Bowel chart is required for all patients – especially those at risk of constipation and consider need for a fluid balance chart & fluid balance summary chart <p>Appendix A, B and C - replaced images.</p> <p>Appendix C - reworked.</p> <p>Appendix E - flowchart revised based on options above in Section 10.</p> <p>Approved at SESLHD Drug and Therapeutics Committee.</p>

Appendix A: Abdominal Assessment

Performing an abdominal assessment, prior to treating a patient for constipation, will help you detect health problems in your patients earlier and prevent further complications from developing.

Have the patient lying flat with a single pillow under their head and the abdomen sufficiently exposed prior to performing examination, otherwise risk activation of abdominal muscles and distortion of anatomy which means examination findings may be unreliable.

Inspection

Visually inspect the abdomen for obvious abnormalities e.g. swelling, bruising, ascites, asymmetry, and localised enlargement. A localised enlargement may indicate a hernia, tumour, cysts, bowel obstruction or enlargement of abdominal organs which can contribute to constipation. A distended abdomen may also be a sign of faecal overload. Any abnormalities need to be reported to the medical team prior to proceeding with any nurse-initiated treatment.

Auscultation

Normal bowel sounds are gurgling sounds (usually occurring between 5 and 35 times per minute) that can be heard with the diaphragm of a stethoscope. Ileus is a condition in which there is a lack of intestinal activity. Decreased sounds, such as no sounds for 1 minute, are a sign of decreased gut activity. Bowel sounds may be noticeably decreased after abdominal surgery, abdominal infection, or injury. Absent sounds (no sounds for 5 minutes) are a warning sign of a potential intestinal obstruction, intestinal perforation, ileus, intestinal ischemia or infarction. A decrease or absence of bowel sounds should be reported to the medical team immediately. Reduced (hypoactive) bowel sounds include a reduction in the loudness, tone, or regularity of the sounds. They are a sign that intestinal activity has slowed. Hypoactive bowel sounds are normal during sleep. They also occur normally for a short time after the use of certain medicines and after abdominal surgery. Decreased or absent bowel sounds often indicate constipation. Increased (hyperactive) bowel sounds can sometimes be heard even without a stethoscope. Hyperactive bowel sounds mean there is an increase in intestinal activity. This may happen with diarrhoea or after eating.

Percussion

With your patient lying supine, percuss all four quadrants of the abdomen using proper technique as follows - hyperextend the middle finger of your non dominant hand and place this finger firmly against your patient's abdomen. With the end (not the pad) of your dominant middle finger, use a quick flick of your wrist to strike the finger on the abdomen. Categorise what you hear as tympanitic or dull.

- **Tympany** is a high pitch (tinkling) sound that is usually present in most of the abdomen caused by air in the bowel, this is associated with a bowel obstruction
- **Dullness** is a flat sound without echoes; that is detected over solid masses such as organs but also in ascites. An unusual dullness may be a clue to an underlying abdominal mass and should be reported to the medical team.

Palpation

To palpate put the fingers of one hand together and depress skin $\frac{1}{2}$ the depth of the abdomen with your fingertips and make gentle rotating movements. The abdomen should be soft and non-tender. Palpate all four quadrants and take note of masses and areas of resistance and tenderness. Any abnormalities and areas of tenderness should be reported to the medical team.




Image adapted by SESLHD using Adobe Stock Imagery.

Abdominal quadrants

The four abdominal quadrants intersect at the umbilicus (navel) — essentially the midpoint of the abdomen.

- A vertical (midsagittal) line runs down the midline of the body through the xiphoid process (bottom of the sternum) and the pubic symphysis.
- A horizontal (transumbilical) line runs through the umbilicus, usually at the level of the L3–L4 vertebrae.

Appendix B: Digital Rectal Examination (DRE)

DRE examination must be performed by a MO, NP/RN/RM or EN with another member of staff present if indicated. The patient's dignity and privacy are always maintained; including considering the gender of the nurse performing the DRE in the context of the patient's gender, expressed preference and cultural/religious background.

Limitations for Practice

- NP/RN/RM or EN must have had education in performing DRE.
- An NP/RN/RM or EN who has not had education should:
 - Read the procedure in this appendix, and
 - Arrange education by consulting with an RN/RM with expertise in DRE or one of the resource personnel listed on page 1.
- EN must always consult an RN/RM prior to performing DRE to determine appropriateness of the procedure and to formulate a treatment plan. Following consultation, DRE is delegated by an RN/RM to the EN if the procedure is indicated.

Indications for Digital Rectal Examination

- To assess anal tone, sphincter function and sensation, characteristics of rectal wall (dilated or collapsed)
- To assess if faecal matter is present, its amount and consistency
- To assess the need for aperients – particularly in patients who have had no bowel action for ≥ 3 days or in whom faecal impaction is suspected.

Clinical history to be obtained prior to digital rectal examination

- Determine whether there are contraindications to performing a DRE (see points 3 and 4)
- Precautions refer to conditions which are not necessarily contraindications, but special care must be taken due to increased risk of rectal trauma in patients with these conditions - medical consultation may be required prior to attending a DRE in these patients
- The decision to perform a DRE on patients with conditions listed under precautions should be made in consultation with the treating medical team and documented in the clinical notes by the treating MO prior to commencing a DRE

Precautions

A clinical history must be obtained prior to proceeding with DRE refer to [Section 2](#).

- Radiotherapy to the pelvic area within the last 6 weeks – consult the Cancer Care Centre nursing staff or treating Radiation Oncologist prior to performing DRE.
- Recent cytotoxic therapy where the patient may potentially be neutropenic.
- Rectal prolapse (depending on degree and any accompanying ulceration) – consult a member of the treating medical team.
- Recent rectal surgery including anterior resection, rectocele repair, rectopexy, haemorrhoidectomy – consult a member of the treating medical team.
- Recent perineal or anal sphincter repair or injury e.g.: post vaginal birth – consult the treating medical officer.

- Known history of sexual or physical abuse – consult with the treating medical team or mental health professional.
- Ensure no known allergies to any of the equipment (i.e.: latex) or anticipated medication.

Contraindications

- Patient does not consent
- Active inflammatory bowel disease including Crohn's disease, ulcerative colitis, diverticulitis
- Rectal excision, anterior resection or ileal pouch surgery within the last 14 weeks
- Trauma to the anal or rectal area
- Obvious per rectal (PR) bleeding
- Rectal or anal pain of unknown cause.
- Anal or rectal lesion/mass
- Tissue fragility
- Spinal injury at risk of autonomic dysreflexia
- Foreign body in rectum or anus
- Imminent terminal event in palliative patient where the procedure will be of questionable value but potentially painful or distressing.
- Thrombocytopenia i.e. decreased platelets of less than 30.

Preparation for procedure

- Inform and educate the patient about the procedure and ensure patient consents to the procedure
- Explain the expected outcome and advise that the urge to defaecate following administration of aperients may occur with minimal warning.
- Collect equipment
 - Blue sheet x 2.
 - Personal protective equipment (PPE) 2 - 3 pairs gloves, disposable apron, facial protection.
 - Lubricant – 2 sachets.
 - Rectal medication as ordered on medication chart or according to Nurse Initiated Medications CIBR.
- Provide an environment to maximise privacy.
- At any time during the procedure if the patient experiences discomfort or becomes agitated:
 - Ask the patient if they wish to continue.
 - If the patient consents to continue the nurse/midwife may proceed.
 - If the procedure is terminated notify/consult with the treating medical team.
 - Document a complete description of the procedure (including if the procedure is terminated) including a description of the patient's reported pain using a pain scale appropriate for the patient.

Procedure

- Perform hand hygiene
- Don PPE
- Position patient in left lateral position (as per picture below)




Image from Adobe Stock Imagery.

- Put protective blue sheets on bed linen
- Observe perianal area for abnormalities.

If the following are detected the nurse should proceed with caution:

- | | |
|------------------|--------------------------------|
| * Anal lesions | * Faecal matter |
| * Haemorrhoids | * Gaping anus |
| * Anal skin tags | * Abnormal perianal skin |
| * Wounds | * Infestations i.e.: parasites |

- Identify anal opening.
- Lubricate index finger and outside of anus.
- Wipe finger across anus to provoke anal reflex.
- Gently insert index finger with the palm facing down.
- During insertion rotate finger through 180 degrees, feeling the circumference of the anal canal.
- Feel to the top of the anal canal and into the base of the rectum. Note any faeces type and consistency and the presence of any masses which are not faeces.
- If any non-faecal mass is detected terminate the procedure, document findings and report to MO.
- Gently withdraw finger.
- Examine withdrawn finger for blood, mucous and colour of stool.
- Discard contaminated glove and perform hand hygiene.
- EN performing procedure discusses findings and treatment plan with RN/RM.

PROCEDURE NOTE

The double glove method may be used by an NP/RN/RM where the first pair of gloves is removed after DRE and the 2nd clean pair used to administer aperients **if indicated**.

An EN who has passed competency assessment may use this method providing the EN has consulted with the RN/RM regarding what findings would indicate the administration of aperients.

Document the following in the clinical notes in eMR and on the eMR Bowel Chart:

- Assessment findings
 - Presence of stool
 - Consistency of stool if present
 - Sphincter tone
 - Characteristics of rectal wall i.e. dilated, collapsed.
 - Presence of haemorrhoids or fissures.
 - Whether further review is indicated and when (refer to [Appendix E](#))
- Procedure outcome including abandoning the procedure if required.

Appendix C: Procedure for Administration of an Enema

All patients requiring the administration of an Enema must have a DRE performed by a clinician suitably trained in the procedure.

1. Fleet Enema


Specific instructions for the Administration of a Fleet Enema at POWH

- An irrigation / enema cone must be used for the administration of a Fleet enema.
- Note: If a patient is deemed to be a bariatric patient, please contact your educator to be advised on a suitable attachment that can be connected to the nozzle of the Fleet enema instead of the cone. This will ensure the Fleet enema solution is delivered accurately and safely via the rectum due to the patient's body habitus.
- A DRE must be performed at least once by a medical officer, usually at the time of prescribing the Fleet enema.
- It is recommended that two nurses are present during the administration of a Fleet enema

Administration of a Fleet Enema

Equipment

- Single use sachet of lubricant
- Enema
- Underpad such as blue sheet
- Disposable gloves / apron / protective eye wear
- The irrigation / enema cone is a soft silicone cone that gently inserts into the rectum in order to safely administer a Fleet enema.
- The irrigation / enema cone **must** be used for the administration of a Fleet Enema at **POWH**
- The tip of the fleet enema is inserted into the irrigation/enema cone tubing and air expelled prior to use.



Generic Procedure for the Administration of an Enema

1. Identify patient and explain procedure and gain consent
2. Place patient in the left lateral position (as per picture below) with a pad under the buttocks




Image from Adobe Stock Imagery.

3. Tilt the head of the bed slightly downward to prevent leakage.
4. Put on disposable gloves and other personal protective equipment as required.

5. Perform DRE examination as per [Appendix B](#).
6. Lubricate the enema or irrigation / enema cone and gently insert into the anus and administer the contents of the Enema Fluid.
7. **If enema or irrigation / enema cone does not insert freely or patient complains of pain, stop immediately and seek medical advice before continuing.**

.....

Generic Procedure for Post Administration of an Enema

8. Remove the enema or irrigation / enema cone and clean anus and buttocks
9. Have pan or commode nearby for patient to use or assist them to the toilet.
10. Instruct the patient to retain the contents of the enema for as long as possible (aim for 15-20 minutes).
11. Document the amount and type of enema and the results in the eMR bowel chart
12. Any complaints of abdominal pain, rectal pain, rectal bleeding, back pain or fever must be promptly and thoroughly evaluated
13. If large volume of faecal output occurs consider possibility of electrolyte imbalance

2. Olive Oil Enema

Olive oil (100-150mL) is retained in the rectum overnight to soften hard or impacted faeces to allow easier evacuation. Do not expect immediate results. The enema may need to be repeated daily before evacuation is achieved. Efficacy depends on the patient's ability to retain the oil. This is usually administered in conjunction with oral aperients or followed by stimulant suppository 12 hours after oil administration.

Preparation of an Olive Oil Enema

- Pour prescribed amount of fluid into an enema bag
- Place enema bag into a bowl of warm water to heat to a tepid temperature.
- Prime tubing to remove air

Perform the Generic Procedure for the Administration of an Enema, Steps 1-7

- Insert rectal catheter of the enema bag in an upward and backward direction for up to 10cm and hold the catheter in position
- **If catheter does not insert freely or patient complains of pain, stop immediately. A mild sensation of the tube being inserted and the enema fluid flowing is expected. Seek medical advice before continuing**
- Attach enema bag to IV pole and allow fluid to slowly flow in.
- Carefully monitor how much fluid the patient can tolerate, usually up to 100 - 150mL can be safely administered; terminate procedure if the patient experiences pain and/or cramps
- If the patient feels like expelling the solution reduce or stop the flow. After the desire to expel the solution has subsided, continue to allow the solution to flow in.

Perform the Generic Procedure for Post Administration of an Enema, Steps 8-13

Appendix D: Procedure for Manual Disimpaction

- Manual disimpaction is not routinely recommended and is used only when other methods of relieving severe constipation have failed. It should only be carried out in consultation with an experienced medical officer.
- Appropriate caution must be used in elderly patients and ensure that consent is taken from substitute decision maker if patient is not able to consent due to significant cognitive impairment.
- Manual disimpaction is not the designated responsibility of either the MO or NP/RN/RM but rather the most experienced clinician.
- Patients are not to be asked to perform self-disimpaction.
- Consult the Palliative Care Team for palliative patients who may require manual faecal disimpaction.
- **SSEH** – Manual disimpaction **must** be performed by a MO.
- Consider early referral to a general surgical team for patients who experience discomfort which would prevent this procedure being carried out on the ward.

Procedure

- 1) Patient may need a mild sedative pre procedure – as per medical team
- 2) Insert 30mL – 60mL of olive oil rectally via rectal tube if possible
- 3) Use 1 finger to gently digitally disimpact rectum so as not to tear or damage the rectal wall
- 4) Clear remaining colon with follow up olive oil enema if required
- 5) Prevent reoccurrence by titrating regular aperients /establishing appropriate bowel management regime and monitoring fluid/food intake

Appendix E: Interventions Related to DRE Findings

