

WATER IMMERSION FOR LABOUR AND BIRTH

This LOP is developed to guide clinical practice at the Royal Hospital for Women. Individual patient circumstances may mean that practice diverges from this LOP.

1. AIM

- To provide woman and staff with appropriate equipment, support, and knowledge about labour and/or birth in water

2. PATIENT

- Woman in labour at term with the following criteria:
 - Singleton pregnancy
 - >37 weeks' gestation
 - Cephalic presentation
 - Normal fetal and maternal observations
 - Booking body mass index (BMI) <35

3. STAFF

- Medical and midwifery staff

4. EQUIPMENT

- Birth pool or appropriate-sized bathtub
- Sieve
- Mirror
- Light/torch
- Gauntlet gloves
- Doppler (waterproof)
- Bath net/sling

5. CLINICAL PRACTICE

- Discuss antenatally with woman the use and benefits of water during first and second stages of labour

WATER IMMERSION FOR LABOUR AND BIRTH - cont'd

- Assess woman's suitability for labouring and/or birthing in water from table 1

Table 1 – Exclusion/Contraindications for Water Immersion:

First Stage	Second Stage
Inability to maintain accurate interpretable continuous electronic fetal monitoring (CEFM) when indicated (as outlined in fetal heart rate monitoring GL2018_025)	
Preterm < 37weeks	Previous shoulder dystocia
Impaired mobility that restricts ability to get in and out of bath	Previous third stage of labour complication e.g. postpartum haemorrhage >1000mls
Abnormal fetal heart rate	Previous obstetric anal sphincter injury (OASI)/severe perineal trauma or history of perineal scarring or fissures (individualised plan needs to be made)
Meconium stained liquor	Known fetal anomaly (individualised plan needs to be made)
Maternal temperature > 37.5°C	Suspected macrosomia
Active herpes simplex infection – Type 1 and 2 Or any open lesion or untreated infection that could impact birth	Known intrauterine growth restriction
Hepatitis B surface antigen positive, Hepatitis C PCR positive or HIV positive	
Vaginal bleeding suggestive of antepartum/intrapartum haemorrhage	
Use of regional anaesthesia	
Opioids within three hours of administration	
Epilepsy or pre-pregnancy diabetes	

WATER IMMERSION FOR LABOUR AND BIRTH – cont'd

- Discuss and refer woman for consultation with senior medical officer who meets the exclusion criteria and is requesting water immersion during first or second stage. There should be a clearly documented individualised care plan as negotiated with the senior obstetric medical officer in the medical record
- Ensure the woman who is eligible for *first stage only* in the water is aware she is advised to exit the bath for second stage
- Ensure woman is aware of circumstances when water immersion is no longer recommended for labour and/or birth (e.g. inability to accurately monitor the fetal heart)
- Ensure water birth is supervised by a midwife experienced with water birth
- Attend routine maternal and fetal observations and record on partogram in line with usual labour and birth care
- Ensure maternal temperature performed hourly and blood pressure (BP) second hourly
- Ensure Intermittent Auscultation (IA), or continuous telemetric cardiotocograph (CTG) are classified as normal. The CTG should be sighted and signed every 15 minutes (as outlined in Fetal Heart Rate Monitoring - Maternity guideline GL2018/025)
- Adjust water temperature according to maternal comfort, aiming to maintain between 35-37.5°C
- Ensure woman adequately hydrated and voiding bladder regularly
- Avoid:
 - soaps, salts, lotions, and oils in water
 - debris – keep water as clear as possible using sieve as much as needed (empty and refill if required)
 - Transcutaneous Electrical Nerve Stimulation (TENS) machine in water
- Ensure neonate is born fully submerged and gently brought to the surface within 10 seconds
- Continue birth out of the water if the birthed head is already exposed to air. Do not re-submerge
- Keep neonate skin-to-skin with woman, keeping warm with blankets and hat whilst maintaining the body submerged. Head must remain above water post birth
- Ensure no tension on the cord to avoid the cord snapping
- Review neonate's condition whilst on woman's chest. If neonatal respirations do not establish or other concern arise, cut the umbilical cord, and transfer the neonate to a resuscitaire for assessment and call for assistance
- Recommend woman exits the bath for active third stage. A woman requesting physiological third stage after a spontaneous labour who wishes to remain in the water should be advised to exit the water if any concern about blood loss arises (see appendix 1 for visual blood loss estimation chart)
- Manage postpartum haemorrhage as outlined in RHW LOP
- Recommend woman leaves pool if concern for her or her neonate's condition in the immediate postpartum period
- Use bath sling if woman is unable to exit bath by herself. Bath sling is located in draw next to bath in birth centre or birth pool trolley in birth unit
- Ensure a minimum of four staff for correct use of the sling
- Do not empty bath as it is easier to get the woman out if bath remains full

WATER IMMERSION FOR LABOUR AND BIRTH – cont'd

Fourth stage of labour

- Recommended that the woman exits the bath once third stage is complete, if has remained in the bath for third stage
- Dry the woman and neonate and keep them warm to prevent hypothermia, keeping the neonate's head dry and covered with a hat, maintaining skin-to-skin as much as possible
- Wrap baby in warmed blankets and give to support person whilst assisting woman to transfer to bed and cover with warmed blankets
- Continue skin-to-skin with neonate
- Delay repair of perineal trauma for an hour (if possible) to allow for skin turgor to return to normal

Shoulder dystocia

- Call for assistance if shoulder dystocia suspected i.e. if the shoulder does not deliver with the next contraction after the neonatal head is born
- Ask the woman to stand up supporting herself with the rails
- Place one of the woman's legs on the side of the bath adopting an upright semi-squat or a deep squat position, if the water level is too high assist the woman out of the bath
- Remove the woman from the bath to the floor if the shoulder is not born with the next contraction and follow the shoulder dystocia procedure according to RHW LOP

6. DOCUMENTATION

Medical record

7. EDUCATIONAL NOTES

- Women reported increased satisfaction with second stage when using water immersion^{1,9}
- Water immersion during the first stage of labour was associated with a small reduction in the risk of using regional analgesia from 43% to 39% (RR 0.91, 95% CI 0.83 to 0.99, 5 trials, 2439 women, moderate-quality evidence)²
- The immersion of a fetal scalp electrode (FSE) in water is currently not supported by the manufacturer – however no adverse outcomes have been observed or documented by them. Therefore, we cannot recommend that immersion in water be used when an FSE is required for fetal monitoring
- There is no evidence of increased maternal, fetal, or neonatal risk associated with water immersion, compared with labouring and giving birth on land^{2,8}. Johnson's review of the newborn respiratory physiology outlines that there are several protective mechanisms that prevent the baby from inhaling or gasping during a birth in water^{7,8}
- There is currently no reliable evidence that can be used to inform women regarding the benefits and risks of warm water immersion during the third stage of labour³ therefore women's wishes and clinical situation should be used to make decision. Following physiological birth, there is no evidence to suggest physiological third stage must be conducted out of the water³. It is important to remember the woman may not show signs of physical compromise until significant blood loss has occurred
- Neonates born in water to Group B Streptococcus (GBS) positive mothers are less frequently colonised with GBS than those born on land⁴

WATER IMMERSION FOR LABOUR AND BIRTH - cont'd

- Women's experience of labour and birth in water is overall very positive. In a physical sense, facilitating ease of movement and assisting women to find a comfortable position as well as the sense of weightlessness being considered helpful. Women suggested that water immersion facilitated a better experience particularly with respect to satisfaction, relaxation, comfort, empowerment and control⁹. This appears to be more significant for those who actually birth in water⁵
- Although there is clearly a need for more research, the currently available evidence does not justify discouraging women from choosing immersion in water during labour^{6,9}
- Fetal hyperthermia has been linked with hypoxia and, therefore, it is important for labouring women to avoid becoming febrile. Monitoring maternal and water temperature regularly is recommended. Water temperature should be comfortable for the woman but not exceed 37.5°C⁶
- It is recommended that annual training of sling use for lifting a woman out of the bath is undertaken by all birthing services staff

8. RELATED POLICIES / PROCEDURES / CLINICAL PRACTICE LOP

- Shoulder Dystocia
- Postpartum Haemorrhage – Prevention and Management
- Environmental Cleaning Policy NSW Health (2020) PD2012_061
- Fetal Heart Rate Monitoring – Maternity – MoH GL2018/025
- Infection Prevention and Control Practice Handbook. Clinical Excellence Commission 2020 NSW

9. RISK RATING

- Low

10. NATIONAL STANDARD

- Standard 5 – Comprehensive Care
- Standard 2 – Partnering with Consumers

11. REFERENCES

1. Pairman S, Pincombe J, Thorogood C, and Tracy S. Midwifery; Preparation for Practice 2015 Churchill Livingstone, Sydney
2. Cluett E, Burns E. 2018. Immersion in water in labour and birth (Review), Cochrane Database of Systematic Reviews
https://www.cochrane.org/CD000111/PREG_immersion-water-labour-and-birth
3. RANZCOG statement March 2021. Warm water immersion during labour and birth. (C-Obs-24) [https://ranzcoq.edu.au/RANZCOG_SITE/media/RANZCOG-MEDIA/Women%27s%20Health/Statement%20and%20guidelines/Clinical-Obstetrics/Water-immersion-during-labour-and-birth-\(C-Obs-24\)_March-2021.pdf?ext=.pdf](https://ranzcoq.edu.au/RANZCOG_SITE/media/RANZCOG-MEDIA/Women%27s%20Health/Statement%20and%20guidelines/Clinical-Obstetrics/Water-immersion-during-labour-and-birth-(C-Obs-24)_March-2021.pdf?ext=.pdf)

WATER IMMERSION FOR LABOUR AND BIRTH - cont'd

4. Zanetti-Dällenbach, R. Lapaire, O., Holzgreve, W. and Hoesli, I. 2007. Neonatal colonization-rate with group B streptococcus is lower in neonates born underwater than after conventional vaginal delivery. *Geburtsh Frauenheilk* 67: 1114–1119
<file:///C:/Users/53041128/Downloads/ZanettiRetal2007.pdf>
5. Cooper, M. and Warland, J. 2019. What are the benefits? Are they concerned? Women's experiences of water immersion for labor and birth. *Midwifery*, vol. 79
<https://www.sciencedirect.com/science/article/abs/pii/S0266613819302335?via%3DiHub>
6. National Institute for Clinical Excellence. Intrapartum care for healthy women and babies Clinical guideline [CG190]. February 2017
7. Johnson P. 1996. Birth under water: To breathe or not to breathe. *BJOG*. 103: 202-8.
8. Harper B. 2014. Birth, Bath, and Beyond: The Science and Safety of Water Immersion During Labor and Birth. *Journal of Perinatal Education* 23(3): 124-34
9. Carlsson T, Ulfsdottir H. 2020. Waterbirth in low-risk pregnancy: An exploration of women's experiences. *Journal of Advanced Nursing*. 76(5): 1221-31
National Institute for Clinical Excellence. Intrapartum care for woman with existing medical conditions or obstetric complication and their babies NICE guideline [NG121]. March, 2019
<https://www.nice.org.uk/guidance/ng121/chapter/Recommendations#previous-caesarean-section>
10. Australian College of Midwives. 2013. Position Statement on the use of water immersion for labour and birth. Australian College of Midwives Canberra

REVISION & APPROVAL HISTORY

Title changed to *Water Immersion for Labour and Birth* Reviewed and endorsed Maternity Services LOPs group 24/8/21

Approved Quality & Patient Safety Committee 21/11/19

Title changed to *Water Immersion for Birth*– reviewed and endorsed Maternity Services LOPs group 5/11/19

Approved Quality & Patient Safety Committee 15/4/11

Reviewed February 2011 and renamed *Waterbirth and Labour in Water*
Labour and Birth in Water approved Quality Council 20/9/04

FOR REVIEW : SEPTEMBER 2026

Visual Estimate of Blood Loss. In an Inflatable Birth Pool



100mls



300mls



500mls



1000mls

Each amount was measured and poured into the water in the inflatable birth pool. Water was 37 degrees Celsius.

Real Blood used (not simulated).