1.In Pae	<u>diatric</u>	Cardio	ic Arre	st ho	w	long	do	you	have	to	pal	pate	for
pulse be	fore st	arting	the ch	est c	om	pres	sio	ns?					

<u>than neonates:</u> <u>Two rescuer:</u> <u>One resucer:</u>
3.If the child is <1 which pulse should be monitored?
4.How do you estimate a child weight based on age?
5.What is the initial energy setting on defibrillator for each of the following: Pulseless VT/VF Unstable SVT Pulsatile VT
6.What are the drug doses for the following resuscitation drugs: Adrenaline IV: Adrenaline I/O Amiodarone: Atropine: HCO3. Suxmaethonium in Neonates:
Child: Adult: Midazolam IV:

- 7. What are the differences between adult and children in insertion technique for Guedel airway?
- 8. What are the different sizes of self inflating bags and for which age group?
- 9. What is estimated calculated ETT depth at the lips for children?
- 10.Until what age Cricothyroid Puncture is the preferred method for oxygenation in difficult airway?
- 11. How much is the initial fluid resuscitation in a child in circulatory shock?

And what is the maximum volume before starting inotrops or blood if hypovolemia is ruled out?

- 12. What is the dose of Pack Cell in children?
- 13. What is the dose for palatelets?
- 14. What is the fluid maintenance for 25 kg child in 24 hours and what type of fluid?

15. What are the likely complication of I/O access line in children?
16.What are the differences between paediatric and adult trauma associated injuries ?
17. How do you roughly calculate ICC size in children?
18.What are the causes of a critically ill neonate?
19.How do you calculate APGAR score?
20.What is the algorithm for neonatal resuscitation?

21.What is the ratio between chest compressions and ventilation in neonates?
22.How many chest compressions and breath you will do in one minute in a neonate?
23.What is the initial O2 Saturation in a term neonate?
24.How many minutes after delivery the O2 sats reading is 85-90%?
25. What is the APGAR Score at 1 min correlates with?
26. APGAR score in 5 min correlates with what?

27.What is the minimal APGAR score that the neonate will
not require any intervention?
28.What is the IV access in order of preference?
30. How many artery and vein does the normal umbilicus
<u>have?</u>
31. How many days after life can umbilical cord be used for
IV access?
32.In emergency situation using umbilical access what type
of catheter and what is the depth of insertion?
of catheter and what is the aepth of miscreton.
33.What are the non-pharmacological pain management?
<u>ooranaa aa o aa o aa aa aa aa aa aa aa aa aa </u>
34.What is the dose of Morphine, Fentanyl in children?

35.What are the key management steps in patients with chronic pain?
36.How many mg present in 1 ml of 10% lignocaine?
37.What is the duration of action of Lignocaine?
38.what is the safe maximum dose of Lignocaine with and without Adrenaline?
39.Regarding local anaethetic agents: Prilocaine safe dose?
Ropivacaine safe dose:
40.Demonstrate:
digital block?

<u>Other blocks</u>
<u>Median nerve</u>
<u>Ulnar nerve</u>
Radial nerve
<u>Tibial nerve</u>
<u>Tibidi liei ve</u>
Fascia iliaca block
41.Intravenous regional anaesthesia indications?
Tiville divollous regional analosomosia maneadistr
Contraindications: