

# Vitamins in cholestasis

## Newborn use only

2025

Alert	The dose recommendations for cholestasis are based on expert opinion. International units (IU) are labelled as units in this formulary. Bio-Logical Vitamin A oral solution and OsteVit D oral liquid contains sodium benzoate. Avoid exposure of >99mg/kg/day in neonates.					
Indication	Neonatal cholestasis					
Action	Vitamin A: Fat soluble vitamin required for vision, growth and bone development, immune function and maintenance of epithelial cells particularly in the retina and respiratory tract tissues. Vitamin D: Regulating levels of calcium and phosphorus and mineralisation of bone. Vitamin E: Antioxidant protecting cell membranes from oxidative stress. Active isomer is α-tocopherol. Vitamin K: Promotes the activation of blood coagulation Factors II, VII, IX and X in the liver.					
Drug type	Fat and water soluble vitamins					
Trade name	Pentavite Infant liquid 0-3 years Brauer Baby Multivitamin Liquid Bio-Logical Vitamin A oral solution Ostelin Vitamin-D3 1000 IU liquid Pretorius Micel-E oral liquid Konakion MM Paediatric					
Presentation	Pentavite Infant – Each 0.45 mL contains 1287 units of vitamin A and 400 units of vitamin D. Brauer Baby Multivitamin Liquid – 1 mL contains 356, 200 and 6 units of vitamin A, D and E. Bio-Logical Vitamin A oral solution – 0.1 mL contains 2500 units of vitamin A. Ostelin Vitamin -D (Vitamin D3) oral liquid – 0.5 mL contains 1000 units of vitamin D. Pretorius Micel-E oral liquid – 0.1 mL contains 15.6 units of vitamin E (equivalent to 10.47 mg) Konakion MM Paediatric – 0.2 mL contains 2 mg vitamin K <sub>1</sub> .					
Dose	Suggested starting regimen <sup>(1-4)</sup>					
		Vitamin A	Vitamin D	Vitamin E	Vitamin K <sub>1</sub>	
	Dose range per day (not per kg)	3000-5000 units	1000-2000 units (25-50 µg)	15-30 units	2 mg twice a week up to 2 mg daily	
	Medical officers to prescribe the following combination of products		Dose Range			
	Vitamins	Oral preparation	Vitamin A (units)	Vitamin D (units)	Vitamin E (units)	Vitamin K <sub>1</sub> (mg)
		Dose (mL) and Frequency				
	Pentavite Infant	0.45 mL once or twice daily	1287-2574	400-800 (10-20 µg)	-	-
	Brauer Baby Multivitamin Liquid-If pentavite is not available	1 mL once or twice daily	356-712	200-400	6-12	
	Bio-Logical Vitamin A solution	0.1 mL daily*	2500	-	-	-

Ostelin Vitamin-D oral liquid 1000 units/0.5 mL <sup>#</sup>	0.5 mL daily*	-	1000 (25 µg)	-	-
Pretorius Micel-E liquid	0.1-0.2 mL daily*	-	-	15.6-31.2	-
Konakion MM Paediatric	0.2 mL twice a week to daily	-	-	-	2 mg twice a week to 2 mg daily

# Vitamins in cholestasis

## Newborn use only

2025

			Total	3787-5074	1400-1800 (35-45 µg)	15.6-31.2	2 mg twice a week to 2 mg daily
	*The daily dose may be administered in two divided doses. #Ostelin Vitamin-D is the preferred liquid because it doesn't contain sodium benzoate. Please check the strengths in each preparation before prescribing.						
Dose adjustment	Not applicable						
Maximum dose							
Total cumulative dose							
Route	Oral or intra-gastric tube.						
Preparation	No preparation is required						
Administration	Administer undiluted or mixed with a small amount of milk into infant's mouth through a feeding teat or via intra-gastric tube.						
Monitoring	Check serum levels of vitamins A, D, E and PT/INR in 1-3 monthly. May need more frequent monitoring in the initial weeks of therapy.						
Contraindications	Hypersensitivity to vitamin A, D, E, K or any component of the formulations. Hypervitaminosis of A, E and/or D.						
Precautions	Direct administration into the mouth may cause choking and apnoea.						
Drug interactions	May increase effects of anticoagulant and antiplatelet agents						
Adverse reactions	Hypervitaminosis A: Irritability, lethargy, vomiting, bulging fontanelle. Hypervitaminosis D: Hypercalcaemia, nephrocalcinosis. Vitamin E: Potentiation of coagulopathy, sepsis, necrotising enterocolitis.						
Overdose	AUSTRALIA: Contact the Poisons Information Centre on 13 11 26 for information on the management of overdose NEW ZEALAND: Contact the National Poisons Centre on 0800 764 766 for information on the management of overdose.						
Compatibility	Not applicable						
Incompatibility	Not applicable						
Stability	Refer to individual product information.						
Storage	All products: Store below 25°C. Protect from light. Pentavite Infant liquid: Refrigerate after opening.						
Excipients	Pentavite Infant liquid: sodium saccharin, pineapple flavour. Brauer Baby Multivitamin Liquid: Ascorbyl palmitate, dl-alpha-tocopherol, gelatin, glycerol, maize oil, maize starch, potassium sorbate, purified water, sucrose, vegetable oil, xanthan gum. Bio-Logical Vitamin A oral solution: sodium benzoate. OsteVit-D oral liquid: sodium benzoate, caramel flavour. Pretorius Micel-E oral liquid: Potassium sorbate and soy bean products. Konakion MM Paediatric: Glycocholic acid, lecithin, sodium hydroxide, hydrochloric acid.						
Special comments	Vitamin E 1 unit = 0.67 mg α-tocopherol. 1 mg of retinyl palmitate = 1818 units of vitamin A.						
Evidence	<b>Background</b> There is a high prevalence of vitamin deficiency in neonatal cholestasis, with one study reporting rates of deficiency for vitamin E (71%), vitamin D (61%), vitamin A (29%) and vitamin K (13%). <sup>(3)</sup> Fat-soluble vitamin serum levels should be monitored frequently in all cholestatic infants to avoid life threatening bleeding (vitamin K deficiency), bone fractures and rickets (vitamin D deficiency), corneal/retinal defects and blindness (vitamin A deficiency), and neurologic and muscular abnormalities (vitamin E deficiency). <sup>(5)</sup> <b>Efficacy</b> Dosing recommendations for vitamins in neonatal cholestasis vary and are based on expert opinion. <sup>(1-4)</sup> Many infants will require individual supplementation of vitamins D, A, E, or K, along with the preferred multivitamin formulation. <sup>(5)</sup> Dosing recommendations for fat soluble vitamins are as follows:						
	Author	Vitamin A	Vitamin D	Vitamin E	Vitamin K		

# Vitamins in cholestasis

## Newborn use only

2025

	Feldman <sup>(4)</sup>	3000-10000 U/day	800-5000 IU/day OR 1,25 OH <sub>2</sub> D <sub>3</sub> : 0.05-0.2 µg/kg/day	Maintain serum targets. No dose recommendations.	2.5- 5 mg twice a week to every day
	Italian society <sup>(1)</sup>	5000-25000 IU/day	800-5000 U/day	15-25 U/kg/day	2.5-5 mg twice a week to every day
	King's college, London <sup>(3)</sup>	1333-5000 IU/day	1000-3000 IU/day	15-150 U/kg/day	1 mg/day
	Lane et al. <sup>(2)</sup>	5000-50000 IU/day	1000-8000 IU/day	1 unit/kg/day	ORAL:2.5-5 mg IM/SQ/IV: 1-10 mg/dose
Optimal approach would be to adjust the doses based on target serum levels. Refer to practice points.					
<b>Safety</b> Excessive doses of vitamins can lead to adverse effects listed in adverse reactions. <sup>(1)</sup>					
<b>Practice points</b>	Aim to maintain the normal range of serum vitamin A, E and D levels. Reference values may vary. Check with your local laboratory. Published recommendations of INR ≤1.2 is often not possible in practice despite high doses of vitamin K. Higher INR values are often accepted as long as there is no clinical evidence of coagulation dysfunction.				
<b>References</b>	<ol style="list-style-type: none"> <li>1. Dani C, Pratesi S, Raimondi F, Romagnoli C. Italian guidelines for the management and treatment of neonatal cholestasis. Italian Journal of Pediatrics. 2015;41:1-12.</li> <li>2. Lane E, Murray KF. Neonatal cholestasis. Pediatric Clinics. 2017;64:621-39.</li> <li>3. Mancell S, Islam M, Dhawan A, Whelan K. Fat-soluble vitamin assessment, deficiency and supplementation in infants with cholestasis. Journal of Human Nutrition and Dietetics. 2022;35:273-9.</li> <li>4. Feldman AG, Sokol RJ. Neonatal cholestasis. Neoreviews. 2013;14(2):e63-e73.</li> <li>5. Feldman AG, Sokol RJ. Neonatal Cholestasis: Updates on Diagnostics, Therapeutics, and Prevention. NeoReviews. 2021;22:e819-e36.</li> </ol>				

VERSION/NUMBER	DATE
Original 1.0	21/07/2022
Version 1.0 (Minor errata)	10/08/2023
Version 1.0 (Minor errata)	28/09/2023
Current 2.0	17/07/2025
REVIEW	17/07/2030

### Authors Contribution of the current version

Author/s	Srinivas Bolisetty, Nilkant Phad
Evidence Review	Srinivas Bolisetty
Expert review of the original	Kate Dehlsen, Scott Nightingale, Michael Stormon, Usha Krishnan, Lizsa Tan, Katie Marks, Andrea Horvath
Nursing Review	Eszter Jozsa, Bryony Malloy
Pharmacy Review	Mohammad Irfan Azeem, Thao Tran
ANMF Group contributors	Bhavesht Mehta, Rebecca Barzegar, Mohammad Irfan Azeem, Rebecca O'Grady, Cindy Chen, Thao Tran, Celia Cunha Brites, Kerrie Knox, Susannah Brew, Bryony Malloy, Renae Gengaroli, Samantha Hassall, Jutta van den Boom, Amber Seigel, Tiffany Kwan, Charles Tian, Emma Watson
Final editing	Srinivas Bolisetty
Electronic version	Thao Tran, Cindy Chen, Helen Huynh, Ian Callander
Facilitator	Srinivas Bolisetty