

Alert	Prolonged ophthalmic use is not recommended due to risk of severe keratitis and corneal adverse effects. For short procedure use only.
Indication	Local anaesthesia for eye examination [including RetCam3™] and procedures [laser] in newborn infants in conjunction with other pharmacological and/or non-pharmacological analgesia methods.
Action	Local anaesthetic.
Drug Type	Ester-type local anaesthetic.
Trade Name	Minims® Oxybuprocaine hydrochloride 0.4% w/v
Presentation	Oxybuprocaine hydrochloride 0.4% single-use, preservative-free, eye drop, approximately 0.5 mL per minim.
Dosage/Interval	One drop each eye as required 1–5 minutes prior to examination. Complete examination within 20 to 30 minutes of administration. Further drops may be needed to achieve a complete anaesthetic effect.
Maximum daily dose	No information.
Route	Topical instillation into the eyes.
Preparation/Dilution	Not applicable.
Administration	Apply pressure to the lacrimal sac during and for 60 seconds after instillation of eye drop to minimise systemic absorption. Wipe away excess medication.
Monitoring	
Contraindications	Hypersensitivity to any of the components of the preparation. Eye infection.
Precautions	The cornea may be damaged by prolonged or frequent application of anaesthetic eye drops. Prolonged use of topical ophthalmic local anaesthetics has been associated with severe keratitis and permanent corneal opacification and scarring with accompanying reduction of visual acuity or visual loss. Systemic toxicity typical of local anaesthetics could occur if sufficient amounts were absorbed systemically. Systemic absorption may be reduced by compressing the lacrimal sac at the medial canthus for a minute during and following the instillation of the drops.
Drug Interactions	Metabolism may be inhibited by anticholinesterases with prolongation of the effects of oxybuprocaine. May competitively enhance the neuromuscular blocking action of suxamethonium.
Adverse Reactions	Transient stinging or burning sensation. Limited safety data in infants. Inappropriate and frequent use might cause corneal ulcerations and perforations.
Compatibility	No information.
Incompatibility	No information.
Stability	Discard immediately after use.
Storage	Store in refrigerator at 2 °C to 8 °C. Do not freeze. Protect from light.
Special Comments	Anaesthesia persists for about 20 to 30 minutes. Corneal sensitivity is normal again after about one hour.
Evidence summary	Efficacy: Eye examination for ROP: No clinical trial has assessed the effect of oxybuprocaine topical eye drops in newborn infants undergoing eye examination for ROP. In an RCT of developmental care measures (NIDCAP) in preterm infants undergoing ROP examination, oxybuprocaine was used in both treatment arms in 20 infants prior to examination.[1, 2] Both treatment arms had significant increases in neonatal pain scores in response to eye examination. [LOE IV] There is more evidence for use of proxymetacaine [proparacaine] eye drops during ROP examination. Three RCTs in newborn infants have evaluated the effects of proxymetacaine hydrochloride 0.5% ophthalmic solution during ROP examination [3–5]. Proxymetacaine eye drops prior to examination were effective in reducing pain responses and/or the number of infants with high pain scores on eye examination. However, the effect was not great and a substantial proportion of infants still had high pain scores.

	<p>Another RCT compared topical anaesthetic eye drops (proxymetacaine hydrochloride 0.5% ophthalmic solution) to oral sucrose during ROP examination. There was no difference between pain scores [6]. A systematic review of these studies concluded that although there is evidence that topical eye anaesthetic at least 30 seconds prior to the procedure reduces pain, its effect is incomplete and screening remains a painful procedure. [7] Conclusion: There is no single pain intervention recommended for ROP examinations [8]. A multimodal approach to eliminating pain is recommended.[2] The Pain Study Group of the Italian Society of Neonatology 2009 recommended, in the case of RetCam3™ screening, applying local anaesthesia with oxybuprocaine 0.4% or tetracaine (amethocaine) 1% eye drops. [9]</p> <p>Laser treatment of ROP: A multicentre, observational study compared local anaesthesia with oxybuprocaine versus intravenous pentazocine versus intravenous fentanyl versus air, oxygen and sevoflurane inhalation.[10] Heart rates were elevated in oxybuprocaine and pentazocine groups; systemic blood pressures were elevated in oxybuprocaine, pentazocine and fentanyl groups; and poor analgesic efficacy was found in the oxybuprocaine, pentazocine and fentanyl groups. However; the air, oxygen and sevoflurane inhalation group experienced hypothermia, enteral feeding intolerance and apnoea more frequently. Conclusion: Oxybuprocaine alone provides inadequate analgesia for laser treatment. [LOE III-2] The Pain Study Group of the Italian Society of Neonatology 2009 recommended, for laser therapy for ROP, in general, combine a local anaesthetic with general anaesthesia. [9]</p> <p>Safety: Older patients describe stinging for 10 seconds after instillation of oxybuprocaine eye drops.[11] There is more evidence for the use of proxymetacaine 0.5% eye drops than oxybuprocaine for ROP screening. However, single use proxymetacaine eye drops are not currently available in Australia and NZ. Proxymetacaine 0.5% is available as 15 mL bottle with preservative benzalkonium chloride. Topical local anaesthetics currently used in ophthalmology, including proxymetacaine and oxybuprocaine, are mostly free of reported complications. Systemic side effects have not been documented with them. However, inappropriate use by patients might cause corneal ulcerations and perforations. Known hypersensitivity to a local anaesthetic drug itself or its preservative is a contraindication for topical local anaesthetic use.[12]</p> <p>Pharmacokinetics: Pharmacokinetics relate to adult data. Oxybuprocaine 0.4% has an intra-ocular surface anaesthesia onset after approximately 1 minute and peak anaesthesia between 1 and 15 minutes. Corneal sensitivity is normal again after about one hour. [13] Anaesthesia persists for about 20 to 30 minutes. Oxybuprocaine is metabolised by plasma and liver esterase enzymes.[13]</p>
<p>References</p>	<ol style="list-style-type: none"> 1. Kleberg A, Warren I, Norman E, Morelius E, Berg AC, Mat-Ali E, Holm K, Fielder A, Nelson N, Hellstrom-Westas L. Lower stress responses after newborn individualized developmental care and assessment program care during eye screening examinations for retinopathy of prematurity: A randomized study. <i>Pediatrics</i>. 2008;121:e1267-e78. 2. Sun X, Lemyre B, Barrowman N, O'Connor M. Pain management during eye examinations for retinopathy of prematurity in preterm infants: A systematic review. <i>Acta Paediatrica, International Journal of Paediatrics</i>. 2010;99:329-34. 3. Cogen MS, Parker JS, Sleep TE, Elsas FJ, Metz Jr TH, McGwin Jr G. Masked trial of topical anesthesia for retinopathy of prematurity eye examinations. <i>Journal of AAPOS</i>. 2011;15:45-8. 4. Marsh VA, Young WO, Dunaway KK, Kissling GE, Carlos RQ, Jones SM, Shockley DH, Weaver NL, Ransom JL, Gal P. Efficacy of topical anesthetics to reduce pain in premature infants during eye examinations for retinopathy of prematurity. <i>Annals of Pharmacotherapy</i>. 2005;39:829-33.

	<p>5. Mehta M, Mansfield T, Vanderveen DK. Effect of topical anesthesia and age on pain scores during retinopathy of prematurity screening. <i>Journal of Perinatology</i>. 2010;30:731-5.</p> <p>6. Nesargi SV, Nithyanandam S, Rao S, Nimbalkar S, Bhat S. Topical anesthesia or oral dextrose for the relief of pain in screening for retinopathy of prematurity: A randomized controlled double-blinded trial. <i>Journal of Tropical Pediatrics</i>. 2015;61:20-4.</p> <p>7. Dempsey E, McCreery K. Local anaesthetic eye drops for prevention of pain in preterm infants undergoing screening for retinopathy of prematurity. <i>Cochrane Database of Systematic Reviews</i> [serial on the Internet]. 2011; (9): Available from: http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD007645.pub2/abstract.</p> <p>8. Francis K. What Is Best Practice for Providing Pain Relief During Retinopathy of Prematurity Eye Examinations? <i>Advances in neonatal care : official journal of the National Association of Neonatal Nurses</i>. 2016;16:220-8.</p> <p>9. Lago P, Garetti E, Merazzi D, Pieragostini L, Ancora G, Pirelli A, Bellieni CV. Guidelines for procedural pain in the newborn. <i>Acta Paediatrica, International Journal of Paediatrics</i>. 2009;98:932-9.</p> <p>10. Sato Y, Oshiro M, Takemoto K, Hosono H, Saito A, Kondo T, Aizu K, Matsusawa M, Futamura Y, Asami T, Terasaki H, Hayakawa M. Multicenter observational study comparing sedation/analgesia protocols for laser photocoagulation treatment of retinopathy of prematurity. <i>Journal of Perinatology</i>. 2015;35:965-9.</p> <p>11. Rossi S et al (Eds). <i>Australian Medicines Handbook 2017</i>. Australian Medicines Handbook Pty Ltd, Adelaide, 2017.</p> <p>12. Gunaydin B, Cok OY. Hazards of topical ophthalmic drug administration. <i>Trends in Anaesthesia and Critical Care</i>. 2011;1:31-4.</p> <p>13. Benoxinate DrugDex Evaluation. DRUGDEX® System (electronic version). Truven Health Analytics. Greenwood Village, Colorado, 2017. Assessed via http://www.micromedexsolutions.com (subscription required) 29/05/17.</p>
--	--

Original version Date: 17/04/2018	Author: NMF Consensus Group
Current Version number: 1.0	Current Version Date: 17/04/2018
Risk Rating: Low	Due for Review: 17/04/2023
Approval by: As per Local policy	Approval Date:

Authors Contribution

Original author/s	Michael Hewson, Cathy Langdon
Expert review	-
Evidence Review	David Osborn
Nursing Review	Eszter Jozsa
Pharmacy Review	Jing Xiao, Cindy Chen
Final content and editing review of the original	Ian Whyte
Electronic version	Cindy Chen, Ian Callander
Facilitator	Srinivas Bolisetty