

# Morphine 10mg/mL (Parenteral)

## Newborn use only

2019

<b>Alert</b>	S8 - High risk medication- may cause significant patient harm when used in error.												
<b>Indication</b>	Analgesia / sedation: 1. Pre-medication prior to intubation or other procedure 2. During assisted ventilation 3. Procedures and post-surgery 4. Neonatal abstinence syndrome secondary to opioid withdrawal												
<b>Action</b>	mu-opioid analgesic – stimulates brain opioid receptors.												
<b>Drug Type</b>	mu-opioid analgesic.												
<b>Trade Name</b>	DBL Morphine Sulfate (also contains sodium chloride and hydrochloric acid). Juno Morphine Hydrochloride												
<b>Presentation</b>	10 mg/mL (10,000 microgram/mL) ampoule. Note: Morphine hydrochloride and sulfate contain approximately equivalent amounts of morphine base per milligram.												
<b>Dosage/Interval</b>	<p><b>ANALGESIA</b></p> <p><b>CONTINUOUS IV INFUSION</b>  Range: 5–40 microgram/kg/hour:  <b>Ventilated infants or after surgery*[1,2,3]</b></p> <table border="1"> <thead> <tr> <th>Postnatal age<sup>#</sup></th> <th>Starting dose</th> <th>Range</th> </tr> </thead> <tbody> <tr> <td>0-7 days</td> <td>10 microgram/kg/hour</td> <td>5-40 microgram/kg/hour</td> </tr> <tr> <td>8-30 days</td> <td>15 microgram/kg/hour</td> <td>5-40 microgram/kg/hour</td> </tr> <tr> <td>31-90 days</td> <td>20 microgram/kg/hour</td> <td>5-40 microgram/kg/hour</td> </tr> </tbody> </table> <p>*Infants after cardiovascular surgery may need lower starting dose and titrated to clinical response.[2]</p> <p><b>IV BOLUS FOR ANALGESIA</b>  50 microgram/kg (maximum recommended 100 microgram/kg) every 4 hours.[4]</p> <p><b>PRE-MEDICATION FOR INTUBATION</b>  100 microgram/kg/dose (up to 200 microgram/kg) [5]</p> <p><b>NEONATAL ABSTINENCE SYNDROME –INITIAL TREATMENT</b>  10 microgram/kg/hour titrated to Neonatal Abstinence Syndrome scores.</p>	Postnatal age <sup>#</sup>	Starting dose	Range	0-7 days	10 microgram/kg/hour	5-40 microgram/kg/hour	8-30 days	15 microgram/kg/hour	5-40 microgram/kg/hour	31-90 days	20 microgram/kg/hour	5-40 microgram/kg/hour
Postnatal age <sup>#</sup>	Starting dose	Range											
0-7 days	10 microgram/kg/hour	5-40 microgram/kg/hour											
8-30 days	15 microgram/kg/hour	5-40 microgram/kg/hour											
31-90 days	20 microgram/kg/hour	5-40 microgram/kg/hour											
<b>Maximum Daily Dose</b>	Doses up to 100 microgram/kg/hour have been used in newborns; however this was associated with an increase in the duration of mechanical ventilation.												
<b>Route</b>	IV												
<b>Preparation/Dilution</b>	<p><b>2-STEP DILUTION (consider for weight &lt;2 kg)</b></p> <p><b>IV Infusion: SINGLE STRENGTH</b></p> <table border="1"> <thead> <tr> <th>Prescribed amount</th> <th>Infusion rate</th> </tr> </thead> <tbody> <tr> <td>1 mg/kg morphine and make up to 50 mL</td> <td>1 mL/hour = 20 microgram/kg/hour</td> </tr> </tbody> </table> <p><b>Step 1:</b> Draw up 1 mL (10mg morphine in 1mL) and add 9 mL sodium chloride 0.9% to make a volume of 10 mL with a concentration of 1000 microgram/mL.  <b>Step 2:</b> From the above solution, draw up 1 mL/kg (1000 microgram/kg) and further dilute with glucose 5% or glucose 10% or sodium chloride 0.9% to make a final volume of 50 mL with a concentration of 1 mL/hour = 20 microgram/kg/hour.  <b>IV bolus dose from single strength solution: 2.5 mL =50 microgram/kg.</b></p> <p><b>IV infusion: DOUBLE STRENGTH</b></p> <table border="1"> <thead> <tr> <th>Prescribed amount</th> <th>Infusion rate</th> </tr> </thead> <tbody> <tr> <td>2 mg/kg morphine and make up to 50 mL</td> <td>1 mL/hour = 40 microgram/kg/hour</td> </tr> </tbody> </table>	Prescribed amount	Infusion rate	1 mg/kg morphine and make up to 50 mL	1 mL/hour = 20 microgram/kg/hour	Prescribed amount	Infusion rate	2 mg/kg morphine and make up to 50 mL	1 mL/hour = 40 microgram/kg/hour				
Prescribed amount	Infusion rate												
1 mg/kg morphine and make up to 50 mL	1 mL/hour = 20 microgram/kg/hour												
Prescribed amount	Infusion rate												
2 mg/kg morphine and make up to 50 mL	1 mL/hour = 40 microgram/kg/hour												

	<p><b>Step 1:</b> Draw up 1 mL (10mg morphine in 1mL) and add 9 mL sodium chloride 0.9% to make a volume of 10 mL with a concentration of 1000 microgram/mL.</p> <p><b>Step 2:</b> From the above solution, draw up 2 mL/kg (2000 microgram/kg) and further dilute with glucose 5% or glucose 10% or sodium chloride 0.9% to make a final volume of 50 mL with a concentration of 1 mL/hour = 40 microgram/kg/hour.</p> <p><b>IV bolus dose from double strength solution: 1.25 mL =50 microgram/kg.</b></p> <p><b>IV BOLUS and PRE-MEDICATION FOR INTUBATION</b>          Draw up 1 mL (10mg morphine in 1mL) and add 9 mL sodium chloride 0.9% to make a final volume of 10mL with a concentration of 1000 microgram/mL.</p> <p><b>1-STEP DILUTION (consider for weight 2 kg and over)</b></p> <p><b>IV Infusion: SINGLE STRENGTH</b></p> <table border="1" data-bbox="472 728 1469 808"> <thead> <tr> <th>Prescribed amount</th> <th>Infusion rate</th> </tr> </thead> <tbody> <tr> <td>1 mg/kg morphine and make up to 50 mL</td> <td>1 mL/hour = 20 microgram/kg/hour</td> </tr> </tbody> </table> <p>Draw up 0.1 mL/kg (10mg morphine in 1mL) and add glucose 5% or glucose 10% or sodium chloride 0.9% to make a final volume of 50 mL with a concentration of 1 mL/hour = 20 microgram/kg/hour.</p> <p><b>For IV bolus dose from single strength solution: 2.5 mL = 50 microgram/kg.</b></p> <p><b>IV Infusion: DOUBLE STRENGTH</b></p> <table border="1" data-bbox="472 1070 1481 1151"> <thead> <tr> <th>Prescribed amount</th> <th>Infusion rate</th> </tr> </thead> <tbody> <tr> <td>2 mg/kg morphine and make up to 50 mL</td> <td>1 mL/hour = 40 microgram/kg/hour</td> </tr> </tbody> </table> <p>Draw up 0.2 mL/kg (10mg morphine in 1mL) and add glucose 5% or glucose 10% or sodium chloride 0.9% to make a final volume of 50 mL with a concentration of 1 mL/hour = 40 microgram/kg/hour.</p> <p><b>For IV bolus dose from double strength solution: 1.25 mL = 50 microgram/kg.</b></p> <p><b>IV BOLUS and PRE-MEDICATION FOR INTUBATION</b>          Draw up 1 mL (10 mg morphine in 1 mL) and add 9 mL sodium chloride 0.9% to make a final volume of 10 mL with a concentration of 1000 microgram/mL.</p>	Prescribed amount	Infusion rate	1 mg/kg morphine and make up to 50 mL	1 mL/hour = 20 microgram/kg/hour	Prescribed amount	Infusion rate	2 mg/kg morphine and make up to 50 mL	1 mL/hour = 40 microgram/kg/hour
Prescribed amount	Infusion rate								
1 mg/kg morphine and make up to 50 mL	1 mL/hour = 20 microgram/kg/hour								
Prescribed amount	Infusion rate								
2 mg/kg morphine and make up to 50 mL	1 mL/hour = 40 microgram/kg/hour								
<b>Administration</b>	<p><b>CONTINUOUS IV INFUSION:</b> Via syringe driver.</p> <p><b>IV BOLUS:</b> Administer over 5 minutes. Flush with 1 mL sodium chloride 0.9% before and after injection. Rapid IV administration may increase adverse effects.</p> <p><b>PRE-MEDICATION FOR INTUBATION:</b> As above for IV bolus. Wait a minimum of 5 minutes for onset of action; however for maximum effect wait 15 minutes after giving the dose.</p>								
<b>Monitoring</b>	<p>All patients should have cardiorespiratory monitoring and be carefully observed, particularly if they are breathing spontaneously. Respiratory depression/apnoea can be reversed with naloxone.</p> <p>Naloxone is contraindicated in opioid dependent infants.</p> <p>Observe for urinary retention, abdominal distension or delay in passage of stool.</p> <p>Withdraw slowly following prolonged use.</p>								
<b>Contraindications</b>	<p>Hypersensitivity to morphine or any excipients.</p>								
<b>Precautions</b>	<p>Potentially toxic serum concentrations of morphine may occur in infants with hypoxic ischaemic encephalopathy with moderate hypothermia and infusion rates &gt;10 microgram/kg per hour. [3] Use with caution in patients with hypersensitivity reactions to other opioids. Hypotension and bradycardia. Respiratory depression.</p>								

# Morphine 10mg/mL (Parenteral)

## Newborn use only

2019

	Transient hypertonia. Convulsions. Ileus and delayed gastric emptying time. Urinary retention. Renal or hepatic impairment. Tolerance may develop after prolonged use – wean slowly.
<b>Drug Interactions</b>	Concomitant use with other CNS depressants potentiates effects of opioids, increasing risk of respiratory depression, profound sedation or coma.
<b>Adverse Reactions</b>	Morphine has been associated with respiratory depression (levels above 20 ng/mL); decreased gastrointestinal motility, hypotension at higher doses, and urinary retention [4].
<b>Compatibility</b>	Compatibility is likely to be similar for morphine hydrochloride and sulfate. <b>Fluids :</b> <b>Morphine hydrochloride</b> – glucose 5%, sodium chloride 0.9% <b>Morphine sulfate</b> – glucose 2.5%, 5% and 10%, glucose in sodium chloride solutions, Hartmann’s, sodium chloride 0.45% and 0.9% <b>Y-site :</b> <b>Morphine hydrochloride</b> –Consult the pharmacist for more advice. <b>Morphine sulfate</b> – adrenaline hydrochloride, amifostine, amikacin, amiodarone, ampicillin, anidulafungin, atracurium, atropine, aztreonam, bivalirudin, caspofungin, cefazolin, cefotaxime, cefoxitin, ceftazidime, ceftriaxone, cisatracurium, clindamycin, dexamethasone, digoxin, dopamine, eptifibatide, erythromycin, esmolol, filgrastim, fluconazole, foscarnet, gentamicin, granisetron, haloperidol lactate (in glucose), heparin sodium, hyoscine hydrobromide, insulin (short-acting), ketorolac, labetalol, lignocaine, linezolid, magnesium sulfate, methylprednisolone sodium succinate, metoclopramide, metoprolol, metronidazole, midazolam, milrinone, noradrenaline, palonosetron, paracetamol, piperacillin-tazobactam (EDTA-free), posaconazole, potassium chloride, remifentanyl, sodium nitroprusside, tacrolimus, tigecycline, tirofiban, tobramycin, trimethoprim-sulfamethoxazole, vancomycin, vecuronium, zidovudine.
<b>Incompatibility</b>	<b>Fluids:</b> Morphine may precipitate out of solution when the final pH is greater than 6.4. <b>Drugs :</b> <b>Morphine hydrochloride</b> – esomeprazole <b>Morphine sulfate</b> – Aminophylline, azathioprine, azithromycin, flucloxacillin, folic acid, ganciclovir, indometacin, pentamidine, pethidine, promethazine, sodium nitrite, thiopental sodium.
<b>Stability</b>	Diluted solution for continuous IV infusion is stable for 48 hours.
<b>Storage</b>	Ampoule: Store below 25°C. Protect from light. Discard remainder after use (in line with schedule 8 drug legislation). Store in Dangerous Drug (DD) safe and record use in DD register.
<b>Special Comments</b>	Prolonged use (> 5–7 days) may be associated with dependence.
<b>Evidence summary</b>	Refer to full version.
<b>References</b>	Refer to full version.

<b>Original version Date: 8/08/2015</b>	<b>Author: Neonatal Medicines Formulary Consensus Group</b>
<b>Current Version number: 2.1</b>	<b>Current Version Date: 19/02/2019</b>
<b>Risk Rating: Medium</b>	<b>Due for Review: 19/02/2022</b>
<b>Approval by: As per Local policy</b>	<b>Approval Date:</b>

### Authors Contribution

Original author/s	Himanshu Popat
Review author/s	Himanshu Popat, Srinivas Bolisetty
Evidence Review	David Osborn
Nursing Review	Eszter Jozsa, Kirsty Minter
Pharmacy Review	Jing Xiao, Cindy Chen, Michelle Jenkins
ANMF Group contributors	Nilkant Phad, Thao Tran
Final content and editing review of the original	Ian Whyte
Electronic version	Cindy Chen, Ian Callander
Facilitator	Srinivas Bolisetty