

Alert	Avoid >5 mg/kg/day as routine supplementation. Check serum ferritin prior to the commencement of medicinal iron following any haemolysis. Consider delaying/temporarily ceasing medicinal iron with (1) multiple transfusions, particularly >100 mL/kg (2) serum ferritin levels >350 microgram/L or (3) have received a transfusion in the last 7 days.																																																												
Indication	<p>1. Prophylaxis in preterm infants <37 weeks and/or birthweight <2.5 Kg 2. Supplementation during erythropoietin therapy 3. Treatment of Iron deficiency anaemia</p> <p>Iron content in the dietary food</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th colspan="3" style="text-align: center;">Iron content</th> </tr> <tr> <th></th> <th style="text-align: center;">140 mL/kg/day</th> <th style="text-align: center;">160 mL/kg/day</th> <th style="text-align: center;">180 mL/kg/day</th> </tr> </thead> <tbody> <tr> <td>Preterm EBM</td> <td style="text-align: center;">0.04 mg/kg/day</td> <td style="text-align: center;">0.05 mg/kg/day</td> <td style="text-align: center;">0.054 mg/kg/day</td> </tr> <tr> <td>EBM+S26 HMF</td> <td style="text-align: center;">0.04 mg/kg/day</td> <td style="text-align: center;">0.05 mg/kg/day</td> <td style="text-align: center;">0.054 mg/kg/day</td> </tr> <tr> <td>EBM+FM 85</td> <td style="text-align: center;">2.1 mg/kg/day</td> <td style="text-align: center;">2.4 mg/kg/day</td> <td style="text-align: center;">2.7 mg/kg/day</td> </tr> <tr> <td>EBM+Nutricia BMF</td> <td style="text-align: center;">0.04 mg/kg/day</td> <td style="text-align: center;">0.05 mg/kg/day</td> <td style="text-align: center;">0.054 mg/kg/day</td> </tr> <tr> <td>Neocate Gold</td> <td style="text-align: center;">1.4 mg/kg/day</td> <td style="text-align: center;">1.6 mg/kg/day</td> <td style="text-align: center;">1.8 mg/kg/day</td> </tr> <tr> <td>Pre Nan Gold</td> <td style="text-align: center;">2.5 mg/kg/day</td> <td style="text-align: center;">2.9 mg/kg/day</td> <td style="text-align: center;">3.2 mg/kg/day</td> </tr> <tr> <td>Aptamil Gold + Preterm</td> <td style="text-align: center;">2.2 mg/kg/day</td> <td style="text-align: center;">2.6 mg/kg/day</td> <td style="text-align: center;">2.9 mg/kg/day</td> </tr> <tr> <td>S26LBW</td> <td style="text-align: center;">2.0 mg/kg/day</td> <td style="text-align: center;">2.2 mg/kg/day</td> <td style="text-align: center;">2.5 mg/kg/day</td> </tr> <tr> <td>Elecare/Elecare LCP</td> <td style="text-align: center;">1.7 mg/kg/day</td> <td style="text-align: center;">1.9 mg/kg/day</td> <td style="text-align: center;">2.2 mg/kg/day</td> </tr> <tr> <td>Pepti-Junior</td> <td style="text-align: center;">1 mg/kg/day</td> <td style="text-align: center;">1.2 mg/kg/day</td> <td style="text-align: center;">1.4 mg/kg/day</td> </tr> <tr> <td>Term Aptamil</td> <td style="text-align: center;">0.78 mg/kg/day</td> <td style="text-align: center;">0.9 mg/kg/day</td> <td style="text-align: center;">1 mg/kg/day</td> </tr> <tr> <td>S26 Gold Newborn</td> <td style="text-align: center;">1.12 mg/kg/day</td> <td style="text-align: center;">1.3 mg/kg/day</td> <td style="text-align: center;">1.4 mg/kg/day</td> </tr> <tr> <td>Nestle NAN Supreme 1</td> <td style="text-align: center;">0.98 mg/kg/day</td> <td style="text-align: center;">1.12 mg/kg/day</td> <td style="text-align: center;">1.26 mg/kg/day</td> </tr> </tbody> </table>		Iron content				140 mL/kg/day	160 mL/kg/day	180 mL/kg/day	Preterm EBM	0.04 mg/kg/day	0.05 mg/kg/day	0.054 mg/kg/day	EBM+S26 HMF	0.04 mg/kg/day	0.05 mg/kg/day	0.054 mg/kg/day	EBM+FM 85	2.1 mg/kg/day	2.4 mg/kg/day	2.7 mg/kg/day	EBM+Nutricia BMF	0.04 mg/kg/day	0.05 mg/kg/day	0.054 mg/kg/day	Neocate Gold	1.4 mg/kg/day	1.6 mg/kg/day	1.8 mg/kg/day	Pre Nan Gold	2.5 mg/kg/day	2.9 mg/kg/day	3.2 mg/kg/day	Aptamil Gold + Preterm	2.2 mg/kg/day	2.6 mg/kg/day	2.9 mg/kg/day	S26LBW	2.0 mg/kg/day	2.2 mg/kg/day	2.5 mg/kg/day	Elecare/Elecare LCP	1.7 mg/kg/day	1.9 mg/kg/day	2.2 mg/kg/day	Pepti-Junior	1 mg/kg/day	1.2 mg/kg/day	1.4 mg/kg/day	Term Aptamil	0.78 mg/kg/day	0.9 mg/kg/day	1 mg/kg/day	S26 Gold Newborn	1.12 mg/kg/day	1.3 mg/kg/day	1.4 mg/kg/day	Nestle NAN Supreme 1	0.98 mg/kg/day	1.12 mg/kg/day	1.26 mg/kg/day
	Iron content																																																												
	140 mL/kg/day	160 mL/kg/day	180 mL/kg/day																																																										
Preterm EBM	0.04 mg/kg/day	0.05 mg/kg/day	0.054 mg/kg/day																																																										
EBM+S26 HMF	0.04 mg/kg/day	0.05 mg/kg/day	0.054 mg/kg/day																																																										
EBM+FM 85	2.1 mg/kg/day	2.4 mg/kg/day	2.7 mg/kg/day																																																										
EBM+Nutricia BMF	0.04 mg/kg/day	0.05 mg/kg/day	0.054 mg/kg/day																																																										
Neocate Gold	1.4 mg/kg/day	1.6 mg/kg/day	1.8 mg/kg/day																																																										
Pre Nan Gold	2.5 mg/kg/day	2.9 mg/kg/day	3.2 mg/kg/day																																																										
Aptamil Gold + Preterm	2.2 mg/kg/day	2.6 mg/kg/day	2.9 mg/kg/day																																																										
S26LBW	2.0 mg/kg/day	2.2 mg/kg/day	2.5 mg/kg/day																																																										
Elecare/Elecare LCP	1.7 mg/kg/day	1.9 mg/kg/day	2.2 mg/kg/day																																																										
Pepti-Junior	1 mg/kg/day	1.2 mg/kg/day	1.4 mg/kg/day																																																										
Term Aptamil	0.78 mg/kg/day	0.9 mg/kg/day	1 mg/kg/day																																																										
S26 Gold Newborn	1.12 mg/kg/day	1.3 mg/kg/day	1.4 mg/kg/day																																																										
Nestle NAN Supreme 1	0.98 mg/kg/day	1.12 mg/kg/day	1.26 mg/kg/day																																																										
Action	Iron is needed for the production of haemoglobin and certain iron-containing enzymes. Ferrous sulphate corrects iron deficiency by re-saturating iron storage organs.																																																												
Drug type	Mineral																																																												
Trade name	ORAL: Ferro-liquid, Maltofer IV – Venofer, Ferrosig iron, Ferrinject																																																												
Presentation	<p>ORAL Ferrous sulfate (ferro-liquid) – 30 mg/mL oral liquid (= 6 mg of elemental iron/mL) Iron polymaltose (Maltofer) – 37 mg/mL (= 10 mg of elemental iron/mL)</p> <p>IV Iron sucrose (Venofer) – 100 mg of elemental iron/5 mL infusion Iron polymaltose (Ferrosig iron) – 100 mg of elemental iron/2 mL Ferric carboxymaltose (Ferrinject) 50 mg of elemental iron/mL injection</p>																																																												
Dose	<p>ORAL</p> <p>1. <u>Iron prophylaxis in preterm infants <37 weeks and/or birthweight <2.5 Kg.</u>⁵⁻⁸ Iron can be from the diet or medicinal iron 2 mg/kg/day - can be commenced from 2 weeks of age and continue up to 6-12 months of age.⁸⁻⁹ Consider delaying/temporarily ceasing iron with (1) multiple transfusions, particularly >100 mL/kg/day, (2) serum ferritin >350 microgram/L or (3) transfusion in the 7 days prior.</p> <p>2. <u>Supplementation during erythropoietin therapy</u> Oral: 3-6 mg/kg/day¹⁰⁻¹¹ IV: 1 mg/kg/day¹²⁻¹³ IV dose of 20 mg/kg/dose can be given weekly¹³</p> <p>3. <u>Treatment of Iron deficiency anaemia</u>⁸ 3-6 mg/kg/day and to continue for 3 months after correction of anaemia⁸</p> <p>IV <u>Supplementation on Parenteral nutrition >4 weeks:</u> Preterm infants: 200-250 micrograms/kg/day¹⁴ or 1400 micrograms/kg weekly¹³ Term infants: 50-100 micrograms/kg/day¹⁴ or 700 micrograms/kg weekly¹³</p> <p><u>Prophylaxis during erythropoietin therapy</u> 1 mg/kg/day¹²</p> <p><u>Treatment of Iron deficiency anaemia:</u> Total Iron dose (mg) = (12.5 – observed Hb (g/dL) x body weight (Kg) x 3.4 x 1.4.¹⁵</p>																																																												

	<p>Oral iron must be ceased 24 hours before IV iron and should not be given until at least 7 days after last parenteral administration.¹⁶</p> <p>IV iron must be prescribed asmg elemental iron (e.g. as iron polymaltose) in.....mL 0.9% sodium chloride over 4 hours.</p> <p>A test dose of 1 mL can be given over 10 minutes prior to the infusion.</p>
Dose adjustment	<p>Therapeutic hypothermia: No information.</p> <p>ECMO: No information.</p> <p>Renal impairment: No information.</p> <p>Hepatic impairment: No information.</p>
Maximum dose	<p>Prophylaxis: 5 mg/kg/day.</p> <p>Treatment: 6 mg/kg/day in iron deficiency anaemia or on erythropoietin.</p>
Total cumulative dose	
Route	<p>ORAL</p> <p>IV</p>
Preparation	<p>ORAL</p> <p>No preparation.</p> <p>IV</p> <p>Draw up required amount of elemental iron from the vial and add to a total volume of sodium chloride 0.9% to a final concentration of 2 mg/mL</p> <p>Example dilution:</p> <p>Total dose of IV iron required is 60mg.</p> <p>Using eg. Ferrosig ampoules containing 100mg elemental iron per 2mL, draw up 1.2mL (60mg) of iron. Add 1.2mL (60mg) to 48.8mL sodium chloride 0.9% to result in a final volume of 50mL with a concentration of 0.024mL (1.2mg) per 1mL.</p>
Administration	<p>ORAL: Administer undiluted.</p> <p>IV: Infusion over 4 hours. A test dose of 1 mL can be given over 10 minutes prior to the infusion.</p>
Monitoring	<p>Periodic haemoglobin and reticulocyte count. Can take 2 weeks for haemoglobin concentrations to rise. Regular serum ferritin if treating iron deficiency anaemia. If the baby has had multiple transfusions, then iron studies would be useful to check for iron overload.</p> <p>IV:</p> <p>Monitor infusion site and for signs of hypersensitivity during and at least for 30 minutes after administration.</p> <p>Continuous cardiorespiratory monitoring, oxygen saturations and temperature.</p>
Contraindications	<p>Anaemia not due to iron deficiency, e.g. chronic haemolytic anaemia</p> <p>Iron overload conditions: haemochromatosis, haemosiderosis</p> <p>Hypersensitivity to iron</p> <p>Uncontrolled hyperparathyroidism</p> <p>Infectious hepatitis – parenteral iron tends to accumulate in inflamed tissues.</p> <p>Acute renal infections – parenteral iron tends to accumulate in inflamed tissues.</p>
Precautions	
Drug interactions	<p>ORAL iron</p> <p>Ascorbic acid favours absorption.</p> <p>Absorption is better if medicinal iron is supplemented with breast milk or between meals; however given with or soon after food may reduce gastrointestinal side effects.¹⁷</p> <p>Not suitable for jejunal administration as enteral absorption occurs in duodenum and upper jejunum. Iron absorption from fortified milk is intact in spite of its high calcium content.</p> <p>IV iron</p> <p>Oral iron is not to be administered concomitantly with IV iron preparations as the absorption of oral iron is reduced. Oral iron therapy should not commence until at least one week after the last iron injection.</p> <p>Concomitant administration of angiotensin converting enzyme (ACE) inhibitors may increase the incidence of adverse effects associated with parental iron preparations eg erythema, abdominal cramps, vomiting and hypotension.</p>
Adverse reactions	<p>ORAL iron</p> <p>GI irritation: abdominal pain, diarrhoea, constipation, dark stools (green or black), gastric mucosal erosion.</p>

	<p>IV iron General: Flushing, sweating, chills and fever; chest and back pain. Hypersensitivity, anaphylaxis. Gastrointestinal: Nausea and vomiting, abdominal pain. Central nervous system: Headache; dizziness. Musculoskeletal: Joint and muscle pain; arthralgia; sensation of stiffening of the arms, legs or face. Cardiovascular: Tachycardia, hypotension, circulatory collapse. Respiratory: Bronchospasm with dyspnoea. Haematological: Generalised lymphadenopathy. Dermatological: Rash, urticarial, angioneurotic oedema. Adverse reactions may be delayed by 1-2 days after treatment with Ferrum H injection.</p> <p>Oral and IV Increased RBC haemolysis and haemolytic anaemia in preterm infants with low vitamin E levels. Rickets - with large doses of iron over a prolonged period of time. Acute toxicity - more severe GI effects including haematemesis and malaena, lethargy, pallor, cyanosis and shock</p>
Compatibility	Can be administered with Penta-vite.
Incompatibility	Do not mix IV solutions with other compounds.
Stability	IV preparations: Venofer: once diluted, use product immediately and discard unused portions. Ferrosig: once diluted, use product immediately and discard unused portions. However if necessary, can store at 2-8°C for not more than 12 hours.
Storage	Store below 25°C. Protect from light
Excipients	Ferro-liquid: sucrose, sorbitol, sodium bisulfite; strawberry flavour Maltofer: ethanol, methyl hydroxybenzoate, propyl hydroxybenzoate, water - purified, sodium hydroxide, sorbitol solution (70%) (non-crystallising), and sucrose. Ferrosig and Ferrum H injections (Iron polymaltose compound): hydrochloric acid or sodium hydroxide (for pH adjustment) Venofer (Iron sucrose): sodium hydroxide (for pH adjustment)
Special comments	Infants on erythropoietin or infants with uncompensated blood loss may initially need higher doses and could be receiving iron supplementation in addition to preterm formula or fortified human milk.
Evidence	Refer to full version.
Practice points	Refer to full version.
References	Refer to full version.

VERSION/NUMBER	DATE
Original	22/10/2015
Revised version 2.0	10/08/2017
Current version 3.0	17/09/2020
REVIEW (5 years)	17/09/2025

Authors Contribution

Original author/s	John Sinn
Current version author/s	Srinivas Bolisetty, John Sinn
Evidence Review	Timothy Schindler
Expert review	Thomas Young, Karel Allegaert
Nursing Review	Eszter Jozsa, Samantha Hassall, Kirsty Minter
Pharmacy Review	Carmen Burman, Mohammad Irfan Azeem
ANMF Group contributors	David Osborn, Nilkant Phad, Bhavesh Mehta, Wendy Huynh, Michelle Jenkins, Thao Tran, Cindy Chen
Final editing and review of the original	Tim Schindler
Electronic version	Cindy Chen, Ian Callander
Facilitator	Srinivas Bolisetty