

Midazolam

Newborn use only

2020

Alert	High risk medication causing significant patient harm when used in error.																
Indication	Sedation during ventilation or procedure. Treatment of refractory seizure.																
Action	Intensify the physiological inhibitory mechanisms mediated by gamma-aminobutyric acid (GABA) by accumulation and occupation of benzodiazepine receptors. Anti-anxiety properties are related to increasing the glycine inhibitory neurotransmitter.																
Drug type	Short acting benzodiazepine.																
Trade name	Hypnovel, Midazolam Alphapharm,, Midazolam Pfizer,, Midazolam-Baxter, B.Braun Midazolam, Midazolam Accord, Midazolam Apotex.																
Presentation	5 mg/mL, 5mg/5mL, 150mg/10mL and 15mg/3mL ampoules used for IV and oral use																
Dose	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 30%;">Method</th> <th>Dose</th> </tr> </thead> <tbody> <tr> <td>IV infusion for sedation</td> <td>0.2–1 microgram/kg/minute</td> </tr> <tr> <td>IV infusion for seizures</td> <td>Loading dose: 150–200 microgram/kg over 3–5 minutes Maintenance dose: 1–7 microgram/kg/minute</td> </tr> <tr> <td>IV bolus</td> <td>50 microgram/kg/dose every 2 hours when required (Dose range: 50–150 microgram/kg/dose)</td> </tr> <tr> <td>IM injection</td> <td>50 microgram/kg/dose every 4 hours when required (Dose range: 50–150 microgram/kg/dose)</td> </tr> <tr> <td>Oral</td> <td>250 microgram/kg as a single dose</td> </tr> <tr> <td>Sublingual</td> <td>200 microgram/kg as a single dose</td> </tr> <tr> <td>Intranasal</td> <td>200 microgram/kg per dose as a single dose (Dose range: 200–300 microgram/kg/dose)</td> </tr> </tbody> </table>	Method	Dose	IV infusion for sedation	0.2–1 microgram/kg/minute	IV infusion for seizures	Loading dose: 150–200 microgram/kg over 3–5 minutes Maintenance dose: 1–7 microgram/kg/minute	IV bolus	50 microgram/kg/dose every 2 hours when required (Dose range: 50–150 microgram/kg/dose)	IM injection	50 microgram/kg/dose every 4 hours when required (Dose range: 50–150 microgram/kg/dose)	Oral	250 microgram/kg as a single dose	Sublingual	200 microgram/kg as a single dose	Intranasal	200 microgram/kg per dose as a single dose (Dose range: 200–300 microgram/kg/dose)
Method	Dose																
IV infusion for sedation	0.2–1 microgram/kg/minute																
IV infusion for seizures	Loading dose: 150–200 microgram/kg over 3–5 minutes Maintenance dose: 1–7 microgram/kg/minute																
IV bolus	50 microgram/kg/dose every 2 hours when required (Dose range: 50–150 microgram/kg/dose)																
IM injection	50 microgram/kg/dose every 4 hours when required (Dose range: 50–150 microgram/kg/dose)																
Oral	250 microgram/kg as a single dose																
Sublingual	200 microgram/kg as a single dose																
Intranasal	200 microgram/kg per dose as a single dose (Dose range: 200–300 microgram/kg/dose)																
Dose adjustment																	
Maximum dose																	
Total cumulative dose																	
Route	IV, IM, Oral, Sublingual, Intranasal.																
Preparation	<p>IV</p> <p>Sedation using 5 mg/1 mL strength</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;"><u>Infusion strength</u></th> <th><u>Prescribed amount</u></th> </tr> </thead> <tbody> <tr> <td><u>1 mL/hour = 1 microgram/kg/minute</u></td> <td><u>3 mg/kg midazolam and make up to 50 mL</u></td> </tr> </tbody> </table> <p>Draw up 0.6 mL/kg (3 mg/kg of midazolam) and add glucose 5%, glucose 10% or sodium chloride 0.9% to make final volume 50 mL. Infuse at a rate of 1 mL/ hour = 1 microgram/kg/minute.</p> <p>Sedation using 5mg/5 mL strength</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;"><u>Infusion strength</u></th> <th><u>Prescribed amount</u></th> </tr> </thead> <tbody> <tr> <td><u>1 mL/hour = 1 microgram/kg/minute</u></td> <td><u>3 mg/kg midazolam and make up to 50 mL</u></td> </tr> </tbody> </table> <p>Draw up 3 mL/kg (3 mg/kg of midazolam) and add glucose 5%, glucose 10% or sodium chloride 0.9% to make final volume 50 mL. Infuse at a rate of 1 mL/ hour = 1 microgram/kg/minute.</p> <p>Seizures using 5 mg/1 mL strength</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;"><u>Infusion strength</u></th> <th><u>Prescribed amount</u></th> </tr> </thead> <tbody> <tr> <td><u>1mL/hour = 5microgram/kg/minutes</u></td> <td><u>15mg/kg midazolam and make up to 50mL</u></td> </tr> </tbody> </table> <p>Draw up 3 mL/kg (15 mg/kg of midazolam) and add glucose 5%, glucose 10% or sodium chloride 0.9% to make final volume 50 mL. Infuse at a rate of 1 mL/hour = 5 microgram/kg/minute.</p> <p>Seizures using 5 mg/5 mL strength (not to be used for babies over 3.3 Kg)</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;"><u>Infusion strength</u></th> <th><u>Prescribed amount</u></th> </tr> </thead> <tbody> <tr> <td></td> <td></td> </tr> </tbody> </table>	<u>Infusion strength</u>	<u>Prescribed amount</u>	<u>1 mL/hour = 1 microgram/kg/minute</u>	<u>3 mg/kg midazolam and make up to 50 mL</u>	<u>Infusion strength</u>	<u>Prescribed amount</u>	<u>1 mL/hour = 1 microgram/kg/minute</u>	<u>3 mg/kg midazolam and make up to 50 mL</u>	<u>Infusion strength</u>	<u>Prescribed amount</u>	<u>1mL/hour = 5microgram/kg/minutes</u>	<u>15mg/kg midazolam and make up to 50mL</u>	<u>Infusion strength</u>	<u>Prescribed amount</u>		
<u>Infusion strength</u>	<u>Prescribed amount</u>																
<u>1 mL/hour = 1 microgram/kg/minute</u>	<u>3 mg/kg midazolam and make up to 50 mL</u>																
<u>Infusion strength</u>	<u>Prescribed amount</u>																
<u>1 mL/hour = 1 microgram/kg/minute</u>	<u>3 mg/kg midazolam and make up to 50 mL</u>																
<u>Infusion strength</u>	<u>Prescribed amount</u>																
<u>1mL/hour = 5microgram/kg/minutes</u>	<u>15mg/kg midazolam and make up to 50mL</u>																
<u>Infusion strength</u>	<u>Prescribed amount</u>																

Midazolam

Newborn use only

2020

	<p><u>1mL/hour = 5microgram/kg/minutes</u></p> <p><u>15mg/kg midazolam and make up to 50mL</u></p> <p>Draw up 15 mL/kg (15 mg/kg of midazolam) and add glucose 5%, glucose 10% or sodium chloride 0.9% to make final volume 50 mL. Infuse at a rate of 1 mL/hour = 5 microgram/kg/minute.</p> <p>IV bolus, IM, oral, sublingual and intranasal</p> <p><u>Using 5 mg/mL ampoule</u>, draw up 0.4 mL (2000 microgram of midazolam) and add 9.6 mL of sodium chloride 0.9% to make final volume of 10 mL with a concentration of 200 microgram/mL.</p> <p><u>Using 5 mg/5mL ampoule</u>, draw up 1 mL (1000 microgram of midazolam) and add 4 mL of sodium chloride 0.9% to make final volume of 5 mL with a concentration of 200 microgram/mL.</p>
Administration	<p>IV infusion: continuous infusion via a syringe pump. Change solution every 24 hours.</p> <p>IV bolus: slow push over 10 minutes.⁹</p> <p>Oral: Plastic IV ampoules may be used for oral administration.</p> <p>Intranasal: IV ampoules may be used for intranasal administration. Drop dose into alternating nostrils over 15 seconds. Absorption is rapid; maximum effect in 10 minutes and duration up to 2 hours. May be irritating to nasal mucosa.</p> <p>IM: Inject deep into a large muscle.</p>
Monitoring	<p>Apnoea, respiratory depression</p> <p>Blood pressure</p> <p>Level of sedation.</p>
Contraindications	<p>Known hypersensitivity to midazolam.</p>
Precautions	<p>In preterm infants, especially in extreme preterm, midazolam half-life is increased from 4–6 hours in term neonates up to 22 hours in premature infants. It is longer with impaired liver function.</p> <p>Caution when concurrently used with opioids – midazolam interacts with other central nervous system depressants and may increase the risk of drowsiness, respiratory depression and hypotension. Withdraw slowly after chronic administration as abrupt discontinuation may precipitate withdrawal seizures.</p> <p>Caution in neonates with renal and hepatic impairment – increased sensitivity to central nervous system (CNS) effects; use doses at lower end of the range.</p> <p>Rapid IV infusion may result in hypotension, respiratory depression or seizure.</p>
Drug interactions	<p>Concurrent administration with erythromycin promotes accumulation.</p> <p>Xanthines may decrease the anaesthetic/sedative effect of benzodiazepines. Care needs to be taken with adding or withdrawing caffeine or aminophylline.</p>
Adverse reactions	<p>Hypotension and reduced cardiac output, particularly when used in combination with fentanyl.</p> <p>Respiratory depression and apnoea.</p> <p>Hypersalivation.</p> <p>Nasal discomfort (with intranasal route).</p> <p>Seizure-like myoclonus (more common in premature neonates receiving via intravenous route).</p>
Compatibility	<p>Fluids: Glucose 5%, glucose 10%, sodium chloride 0.9%.</p> <p>Y-site: Amino acid solutions. Abciximab, amikacin, amiodarone, anidulafungin, atracurium, bivalirudin, calcium gluconate, caspofungin, cefotaxime cefazolin, ciprofloxacin, cisatracurium, clindamycin, digoxin, dopamine, doripenem, eptifibatide, erythromycin, esmolol, fentanyl, fluconazole, gentamicin, glyceryl trinitrate, haloperidol lactate, hydromorphone, labetalol, linezolid, methadone, methylprednisolone, metronidazole, milrinone, morphine sulfate, noradrenaline (norepinephrine), palonosetron, pancuronium, potassium chloride, ranitidine, remifentanyl, sodium nitroprusside, tirofiban, tobramycin, vancomycin, vecuronium.</p>
Incompatibility	<p>Fluids: No information.</p> <p>Y-site: Fat emulsion. Aciclovir, albumin, aminophylline, amoxicillin, ampicillin, azathioprine, azithromycin, cefepime, ceftazidime, chloramphenicol, clonidine, dexamethasone, ertapenem, esomeprazole, flucloxacillin, foscarnet, furosemide (frusemide), ganciclovir, hydrocortisone sodium succinate, imipenem-cilastatin, indomethacin, omeprazole, phenobarbital (phenobarbitone), piperacillin-tazobactam (EDTA-free), potassium acetate, sodium bicarbonate, thiopental (thiopentone), tramadol, trimethoprim-sulfamethoxazole.</p>
Stability	<p>Diluted solution: Store at 2–8°C and use within 24 hours.</p>

Storage	Midazolam Apotex, Midazolam-Baxter: Store below 30°C. Protect from light. B. Braun Midazolam, Hypnovel, Midazolam Alphapharm: Store below 25°C. Protect from light. Midazolam Pfizer: Store below 25°C. Protect from light. Unopened ampoules will be suitable for use for up to 8 months after the foil sachet has been opened, if protected from light. Schedule 4D (S4D) medication. Store in dangerous drug safe and record use in S4D register.
Excipients	Sodium chloride, hydrochloric acid, sodium hydroxide, water for injections.
Special comments	Flumazenil is a specific benzodiazepine antagonist and may be used (very limited experience in the neonate) to rapidly reverse respiratory depression – 10 microgram/kg/dose IV push. May repeat every minute for up to 4 more doses.
Evidence	<p>Efficacy</p> <p>There are insufficient data to promote the use of intravenous midazolam infusion as a sedative for neonates undergoing intensive care. Although all studies included in the review reported better sedation, none of the scales used had been validated in preterm infants and thus the effectiveness could not be evaluated [1] (Level 1, Grade B).</p> <p>Midazolam was effective in neonates with refractory seizures that did not respond to phenobarbital (phenobarbitone), phenytoin or pentobarbital (pentobarbitone) [2] (Level IV, Grade D).</p> <p>Safety</p> <p>One study showed a statistically significant higher incidence of adverse neurological events (death, grade III or IV IVH, PVL) and meta-analysis of data from two studies showed a statistically significant longer duration of NICU stay in the midazolam group compared to the placebo group [1] (Level 1, Grade B).</p> <p>Administration of midazolam in ventilated premature infants causes significant changes in cerebral oxygenation and hemodynamics, which might be harmful [3] (Level III, Grade C).</p> <p>Intravenous bolus doses of midazolam in association with fentanyl should be used with great caution in the newborn, especially if very premature or with unstable blood pressure [4] (Level IV, Grade D).</p> <p>Sedation with midazolam has a transient effect on the background aEEG activity [5] (Level III, Grade C).</p> <p>Pharmacokinetics</p> <p>Midazolam is highly protein bound with an elimination half-life of 4–6 hours in term neonates and a variable half-life (up to 22 hours) in premature neonates and those with impaired hepatic function. Bioavailability is approximately 36% with oral administration and 50% with sublingual and intranasal administration [6] (Level III, Grade C).</p>
Practice points	
References	<ol style="list-style-type: none"> 1. Ng E, Taddio A, Ohlsson A. Intravenous midazolam infusion for sedation of infants in the neonatal intensive care unit. <i>The Cochrane database of systematic reviews</i>. 2012;6:CD002052. 2. Castro Conde JR, Hernandez Borges AA, Domenech Martinez E, Gonzalez Campo C, Perera Soler R. Midazolam in neonatal seizures with no response to phenobarbital. <i>Neurology</i>. Mar 8 2005;64(5):876–879. 3. Van Alfen-van der Velden AA, Hopman JC, Klaessens JH, Feuth T, Sengers RC, Liem KD. Effects of midazolam and morphine on cerebral oxygenation and hemodynamics in ventilated premature infants. <i>Biology of the Neonate</i>. 2006;90(3):197–202. 4. Burtin P, Daoud P, Jacqz-Aigrain E, Mussat P, Moriette G. Hypotension with midazolam and fentanyl in the newborn. <i>Lancet</i>. Jun 22 1991;337(8756):1545–1546 5. Bernet V, Latal B, Natalucci G, Doell C, Ziegler A, Wohlrab G. Effect of sedation and analgesia on postoperative amplitude-integrated EEG in newborn cardiac patients. <i>Pediatr Res</i>. Jun 2010;67(6):650–655. 6. De Wildt SN, Kearns GL, Hop WC, Murry DJ, Abdel-Rahman SM, van den Anker JN. Pharmacokinetics and metabolism of intravenous midazolam in preterm infants. <i>Clin Pharmacol Ther</i>. 2001 Dec;70(6):525–31. 7. Taketomo CK, Hodding JH, Kraus DM, American Pharmacists Association. <i>Pediatric and neonatal dosage handbook</i>. Hudson, Ohio: Lexi-Comp: American Pharmacists Association; 2015. 8. <i>Australian Injectable Drugs Handbook, 6th Edition, Society of Hospital Pharmacists of Australia 2014.</i>

	9. Van Den Broek MP, Van Straaten HL, Huitema AD, Egberts T, Toet MC, De Vries LS, Rademaker K, Groenendaal F. Anticonvulsant effectiveness and hemodynamic safety of midazolam in full-term infants treated with hypothermia. <i>Neonatology</i> . 2015 Jan 8;107(2):150-6.
--	--

VERSION/NUMBER	DATE
Original 1.2	23/03/2016
Version 2.0	16/11/2020
REVIEW (5 years)	16/11/2025

Authors Contribution

Original author/s	David Osborn, Srinivas Bolisetty
Evidence Review	David Osborn
Expert review	
Nursing Review	Eszter Jozsa, Kirsty Minter
Pharmacy Review	Thao Tran, Michelle Jenkins
ANMF Group contributors	Himanshu Popat, Chris Wake, Nilkant Phad, Bhavesh Mehta, John Sinn, Carmen Burman, Jessica Mehegan, Wendy Huynh, Helen Huynh
Final editing and review of the original	Ian Whyte
Electronic version	Cindy Chen, Ian Callander
Facilitator	Srinivas Bolisetty