

Zidovudine

Newborn use only

2020

Alert	No Australian registered intravenous products are available. Retrovir IV ampoules are only available via the Special Access Scheme (SAS) in Australia. Also known as azidothymidine (AZT).																					
Indication	Monotherapy or part of a combination therapy for prevention of maternal-foetal HIV transmission.																					
Action	Nucleoside analogue that inhibits HIV replication by interfering with viral reverse transcriptase.																					
Drug type	Antiretroviral medication																					
Trade name	Retrovir																					
Presentation	Oral: syrup 10 mg/mL IV: 10 mg/mL in a 20mL single-use vial (SAS) Note: Retrovir is also available in oral capsules, however only the syrup is used in neonates.																					
Dose	<p>Oral Start therapy within 4 hours of birth.</p> <table border="1"> <thead> <tr> <th>Gestation at birth</th> <th>Dose</th> <th>Interval</th> </tr> </thead> <tbody> <tr> <td><30 weeks</td> <td>2 mg/kg</td> <td>12 hourly</td> </tr> <tr> <td>30⁺⁰-33⁺⁶ weeks</td> <td>2 mg/kg</td> <td>12 hourly for 2 weeks and then 8 hourly</td> </tr> <tr> <td>≥34 weeks</td> <td>4 mg/kg</td> <td>12 hourly*</td> </tr> </tbody> </table> <p>*Dose can be rounded up to the nearest 0.5 mg to assist administration.</p> <p>IV If neonates are unable to take oral zidovudine</p> <table border="1"> <thead> <tr> <th>Gestation at birth</th> <th>Dose</th> <th>Interval</th> </tr> </thead> <tbody> <tr> <td>≤33⁺⁶ weeks gestation*</td> <td>1.5 mg/kg/dose</td> <td>12 hourly</td> </tr> <tr> <td>≥34 weeks gestation</td> <td>1.5 mg/kg/dose</td> <td>6 hourly</td> </tr> </tbody> </table> <p>* Change interval to 6 hourly at 34 weeks gestation. Switch to oral once the neonate is tolerating oral feeds.</p> <p>Total duration IV/oral dosing</p> <ul style="list-style-type: none"> • Very low risk monotherapy – 2 weeks • Low risk monotherapy – 4 weeks • High risk / combination therapy – 4 weeks 	Gestation at birth	Dose	Interval	<30 weeks	2 mg/kg	12 hourly	30 ⁺⁰ -33 ⁺⁶ weeks	2 mg/kg	12 hourly for 2 weeks and then 8 hourly	≥34 weeks	4 mg/kg	12 hourly*	Gestation at birth	Dose	Interval	≤33 ⁺⁶ weeks gestation*	1.5 mg/kg/dose	12 hourly	≥34 weeks gestation	1.5 mg/kg/dose	6 hourly
Gestation at birth	Dose	Interval																				
<30 weeks	2 mg/kg	12 hourly																				
30 ⁺⁰ -33 ⁺⁶ weeks	2 mg/kg	12 hourly for 2 weeks and then 8 hourly																				
≥34 weeks	4 mg/kg	12 hourly*																				
Gestation at birth	Dose	Interval																				
≤33 ⁺⁶ weeks gestation*	1.5 mg/kg/dose	12 hourly																				
≥34 weeks gestation	1.5 mg/kg/dose	6 hourly																				
Dose adjustment	Therapeutic hypothermia: no information. ECMO: no information. Renal: see monitoring and interactions. Hepatic: see monitoring and adverse reactions.																					
Maximum dose																						
Total cumulative dose																						
Route	Oral IV																					
Preparation	Oral: Syrup IV: Draw up 1mL (10mg of zidovudine) and add 9mL of glucose 5% to make a final volume of 10mL with a final concentration of 1mg/mL. [1]																					
Administration	Oral: Can be given without food. IV: infusion over 30 minutes - 1 hour.																					
Monitoring	Full blood count, blood sugar level, liver function, renal functions, viral load, CD4 counts should be obtained. The panel should be repeated within 2-4 weeks of commencement of therapy and then every 3-4 months. [2-4]																					
Contraindications	Life-threatening hypersensitivity reactions (e.g., anaphylaxis, Stevens-Johnson syndrome) to zidovudine or any components of the formulations. [5] Zidovudine infusions should not be given to patients with abnormally low neutrophils or haemoglobin levels. [5]																					

Zidovudine

Newborn use only

2020

Precautions	There have been reports of pancytopenia associated with the use of zidovudine, which was reversible in most instances after discontinuance of the drug.
Drug interactions	<p>Stavudine - zidovudine should not be administered in combination with stavudine because of in vitro virologic antagonism.</p> <p>Co-administration of zidovudine with drugs that are nephrotoxic, cytotoxic, or which interfere with red blood cell and white blood cell number or function (e.g. ganciclovir, amphotericin B or interferon) may increase the risk of toxicity. If concomitant therapy with any of these drugs is necessary then extra care should be taken in monitoring renal function and haematological parameters.</p> <p>Ribavirin antagonizes in vitro antiviral activity of zidovudine and so concomitant use should be avoided.</p> <p>Doxorubicin - simultaneous use of doxorubicin and zidovudine should be avoided. Doxorubicin may inhibit the phosphorylation of zidovudine to its active form.</p> <p>Phenytoin - phenytoin blood levels have been reported to be low in some patients receiving zidovudine. Monitor phenytoin levels if neonate is receiving both medications. [5]</p> <p>Clarithromycin - oral clarithromycin reduces the absorption of zidovudine. This can be avoided by separating the doses by at least 2 hours. [5]</p>
Adverse reactions	Anaemia and neutropenia are common. Transient lactic acidemia, vomiting, headache, insomnia, hepatomegaly with hepatic steatosis, lipodystrophy, lipoatrophy, myopathy, cardiomyopathy and myositis. [6, 7] In most cases the adverse events are mild and self-limiting. Prolonged use increases the risk of adverse events.
Compatibility	<p>Fluids: glucose 5%, sodium chloride 0.9%</p> <p>Y site: aciclovir, amikacin, amphotericin B, aztreonam, cefepime, ceftazidime, ceftriaxone, cimetidine, clindamycin, dexamethasone, dobutamine, dopamine, erythromycin lactobionate, fluconazole, gentamicin, heparin, imipenem, linezolid, lorazepam, metoclopramide, morphine, nafcillin, oxacillin, piperacillin, piperacillin-tazobactam, potassium chloride, ranitidine, remifentanyl, rocuronium, tobramycin, trimethoprim-sulfamethoxazole, and vancomycin.</p> <p>Note: This is not an exhaustive list. Please refer to the relevant resources eg. Micromedex, Australian Injectable Drugs Handbook for detailed information.</p>
Incompatibility	<p>Fluids: no information</p> <p>Y site: lansoprazole, meropenem</p>
Stability	<p>Vial: store below 30°C</p> <p>After dilution, the drug solution is stable for 24 hours if stored below 25°C or in refrigerator. Protect from light. [5]</p>
Storage	Oral syrup and any unused vials are to be stored at room temperature and protected from light. [5]
Excipients	<p>Retrovir Oral Syrup: Each 5 mL contains zidovudine 50 mg, and glycerol, citric acid, sodium benzoate, saccharin sodium, maltitol solution, Flavour Strawberry 500286E, Flavour White Sugar 3112044, and water-purified.</p> <p>Retrovir IV vials: hydrochloric acid, sodium hydroxide, water for injection.</p>
Special comments	<p>Dose adjustment is required in renal and hepatic impairment.</p> <p>Fixed drug combinations should be avoided in infants with renal and hepatic insufficiency.</p>
Evidence	Refer to full version.
Practice points	Refer to full version.
References	Refer to full version.

VERSION/NUMBER	DATE
Original 1.0	29/05/2020
Current 1.1	16/11/2020
REVIEW	16/11/2025

Authors Contribution

Original author/s	Nilkant Phad, Srinivas Bolisetty
Evidence Review	David Osborn

Expert review	Pamela Palasanthiran
Nursing Review	Eszter Jozsa, Kirsty Minter
Pharmacy Review	Carmen Burman, Cindy Chen, Michelle Jenkins
ANMF Group contributors	Himanshu Papat, Carmen Burman, Thao Tran, John Sinn, Emily Do, Wendy Huynh, Bhavesh Mehta, Rahul Udaya Prasad
Final editing and review of the original	David Osborn, Michelle Jenkins, Srinivas Bolisetty
Electronic version	Ian Callander, Cindy Chen
Facilitator	Dr Srinivas Bolisetty