

CLINICAL POLICIES AND PROCEDURES

**NEWBORN USE ONLY
GIVEN ON DOCTORS ORDER ONLY**

INTUBATION AND EMERGENCY DRUGS

INTUBATION DRUGS	Dose	Dilution
Morphine <i>10mg / 1 ml</i>	100 mcg / kg	9 ml WFI + 1 ml Morphine = 1mg/1ml
Atropine <i>600mcg / 1 ml</i>	10 mcg / kg	5 ml WFI + 1 ml Atropine =100mcg/1ml
Suxamethonium <i>100mg / 2 ml</i>	1 – 2 mg / kg	8 ml WFI + 2 ml Sux.= 10mg/1ml

EMERGENCY DRUGS	Dose	Dilution
Adrenaline 1:10 000	0.1 ml / kg	None
Atropine <i>600 mcg / 1 ml</i>	10 mcg / kg	5 ml WFI + 1 ml Atropine =100mcg/1ml
Calcium Gluconate 10% <i>2.2 mmol in 10 ml</i>	1 ml / kg	5ml 5%dex + 5ml Cal. Gluc.=0.11mmol /1ml
Sodium Bicarbonate 8.4% <i>10 mmol in 10 ml</i>	1 – 3 mmol / kg	5 ml WFI + 5 ml Sodi. Bic. 2 ml = 1 mmol
Naloxone <i>400 mcg / 1 ml</i>	100 mcg / kg	None